Native Harvest of Wildlife in the Keewatin Region, Northwest Territories for the Period October 1985 to March 1986 and a Summary for the Entire Period of the Harvest Study from October 1981 to March 1986

R.L. Gamble

Central and Arctic Region
Department of Fisheries and Oceans
Winnipeg, Manitoba R3T 2N6

March 1988

Canadian Data Report of Fisheries and Aquatic Sciences No. 688



88-03406/

Canadian Data Report of Fisheries and Aquatic Sciences

These reports provide a medium for filing and archiving data compilations where little or no analysis is included. Such compilations commonly will have been prepared in support of other journal publications or reports. The subject matter of Data Reports reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries management, technology and development, ocean sciences, and aquatic environments relevant to Canada.

Numbers 1-25 in this series were issued as Fisheries and Marine Service Data Records. Numbers 26-160 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Data Reports. The current series name was changed with report number 161.

The correct citation appears above the abstract of each report.

Rapport statistique canadien des sciences halieutiques et aquatiques

Ces rapports servent de base à la compilation des données de classement et d'archives pour lesquelles il y a peu ou point d'analyse. Cette compilation aura d'ordinaire été préparée pour appuyer d'autres publications ou rapports. Les sujets des Rapports statistiques reflètent la vaste gamme des intérêts et politiques du Ministère des Pêches et des Océans, notamment gestion des pêches, techniques et développement, sciences océaniques et environnements aquatiques, au Canada.

Les numéros 1 à 25 de cette série ont été publiés à titre de Records statistiques, Service des pêches et de la mer. Les numéros 26-160 ont été publiés à titre de Rapports statistiques du Service des pêches et de la mer, Ministère des Pêches et de l'Environnement. Le nom de la série a été modifié à partir du numéro 161.

Le titre exact paraît au haut du résumé de chaque rapport.

Canadian Data Report of

Fisheries and Aquatic Sciences 688

MARCH 1988

NATIVE HARVEST OF WILDLIFE IN THE KEEWATIN REGION, NORTHWEST TERRITORIES FOR THE PERIOD OCTOBER 1985 TO MARCH 1986 AND A SUMMARY FOR THE ENTIRE PERIOD OF THE HARVEST STUDY FROM OCTOBER 1981 TO MARCH 1986

by

R. L. Gamble[1]

Central and Arctic Region

Department of Fisheries and Oceans

Winnipeg, Manitoba R3T 2N6

This is the 22nd Data Report

from the Central and Arctic Region, Winnipeg

---

[1] Present address 20 Amundsen Bay, Winnipeg, Manitoba R3K 0V2

Former address Keewatin Wildlife Federation, Rankin Inlet, N.W.T. X0C 0G0

PREFACE

This report is a supplement to previous technical reports by Gamble (1984, 1987a, 1987b) which cover the period October 1981 to September 1985 and covers the period October 1985 to March 1986. It is presented in fulfillment of Department of Supply and Services Contract DSS 25 S.T.A. 7135-05-0003 let to the Keewatin Wildlife Federation to conduct a wildlife harvest study in the Keewatin Region - Phase III. The work was done on behalf of the Federal Government departments of Environment Canada (Canadian Wildlife Service), Fisheries and Oceans (Central and Arctic Region), and Indian Affairs and Northern Development; the Government of the Northwest Territories Department of Renewable Resources; and the Keewatin Wildlife Federation.

The report is accepted upon recommendation by the steering committee for the study made up of representatives of the agencies noted above (Appendix 3). The harvest study material is published under the auspices of the DFO report series by agreement of the steering committee in order to ensure that the data achieve a wide circulation, be accessible to the interested public, and be published in a standardized format generally recognized as appropriate for the dissemination of such information.

A report of the study in Inuktitut will also be published under the auspices of Indian Affairs and Northern Development.

© Minister of Supply and Services Canada 1988

Cat. no. Fs 97-13/688E ISSN 0706-6465

Correct citation for this publication is:

Gamble, R.L. 1988. Native harvest of wildlife in the Keewatin Region, Northwest Territories for the period October 1985 to March 1986 and a summary for the entire period of the harvest study from October 1981 to March 1986. Can. Data Rep. Fish. Aquat. Sci. 688: v + 85 p.

TABLE OF CONTENTS

	Page
ABSTRACT/RÉSUMÉ	v
INTRODUCTION	1
MATERIALS AND METHODS	1
General	1
Human resources	1
Materials	1
Data collection and analysis	1
Data processing	2
ACKNOWLEDGMENTS	2
REFERENCES	2

PART I. NATIVE HARVEST FOR THE PERIOD OCTOBER 1985 TO MARCH 1986.

LIST OF TABLES

Table		Page
1	The reported harvest by Baker Lake hunters, for the period October 1985 to March 1986	4
2	The reported harvest by Chesterfield Inlet hunters, for the period October 1985 to March 1986	4
3	The reported harvest by Coral Harbour hunters, for the period October 1985 to March 1986	5
4	The reported harvest by Eskimo Point hunters, for the period October 1985 to March 1986	5
5	The reported harvest by Rankin Inlet hunters, for the period October 1985 to March 1986	6
6	The reported harvest by Repulse Bay hunters, for the period October 1985 to March 1986	6
7	The reported harvest by Whale Cove hunters, for the period October 1985 to March 1986	7
8	The estimated harvest by Baker Lake hunters, for the period October 1985 to March 1986	7
9	The estimated harvest by Chesterfield Inlet hunters, for the period October 1985 to March 1986	8
10	The estimated harvest by Coral Harbour hunters, for the period October 1985 to March 1986	8
11	The estimated harvest by Eskimo Point hunters, for the period October 1985 to March 1986	9
12	The estimated harvest by Rankin Inlet hunters, for the period October 1985 to March 1986	9
13	The estimated harvest by Repulse Bay hunters, for the period October 1985 to March 1986	10
14	The estimated harvest by Whale Cove hunters, for the period October 1985 to March 1986	10
15	The reported and estimated harvest for Baker Lake hunters	11
16	The reported and estimated harvest for Chesterfield Inlet hunters	11
17	The reported and estimated harvest for Coral Harbour hunters	12
18	The reported and estimated harvest for Eskimo Point hunters	12
19	The reported and estimated harvest for Rankin Inlet hunters	13
20	The reported and estimated harvest for Repulse Bay hunters	13
21	The reported and estimated harvest for Whale Cove hunters	14
22	Monthly theoretical kill factors for seven Keewatin communities	14
23	The harvest by species over the range of age for Baker Lake hunters covering the period October 1985 to March 1986	14
24	The harvest by species over the range of age for Chesterfield Inlet hunters covering the period October 1985 to March 1986	15
25	The harvest by species over the range of age for Coral Harbour hunters covering the period October 1985 to March 1986	15
26	The harvest by species over the range of age for Eskimo Point hunters covering the period October 1985 to March 1986	16
27	The harvest by species over the range of age for Rankin Inlet hunters covering the period October 1985 to March 1986	16
28	The harvest by species over the range of age for Repulse Bay hunters covering the period October 1985 to March 1986	17
29	The harvest by species over the range of age for Whale Cove hunters covering the period October 1985 to March 1986	17
30	Data on the distribution of hunters that were successful in obtaining a harvest, for the period October 1985 to March 1986	18
31	Edible weight values in kilograms for harvested species as calculated from various sources	19

Table		Page
32	Reported and estimated edible weight values for harvested species for the period October 1985 to March 1986 .	20
33	Reported and estimated edible weight values for four major groups of animals harvested by Keewatin communities, October 1985 to March 1986 . .	22
34	Age distribution of hunters for the seven Keewatin region communities for the period October 1985 to March 1986.	26

PART II. SUMMARY OF NATIVE HARVEST FOR THE PERIOD OCTOBER 1981 TO MARCH 1986 (APPENDIX 4 GIVES FOOTNOTES).

LIST OF TABLES

Table		Page
35	The estimated harvest by Baker Lake hunters, for the period November 1981 to March 1986	27
36	The estimated harvest by Chesterfield Inlet hunters, for the period January, February and August 1982 through March 1936	33
37	The estimated harvest by Coral Harbour hunters, for the period February 1982 and June 1982 through April 1983 and October 1983 to March 1986 . .	37
38	The estimated harvest by Eskimo Point hunters, for the period October 1981 to March 1986	44
39	The estimated harvest by Rankin Inlet hunters, for the period November 1981 to March 1986	51
40	The estimated harvest by Repulse Bay hunters, for the period October 1981 to March 1986	57
41	The estimated harvest by Whale Cove hunters, for the period October 1981 to March 1983 and November 1983 to March 1986	63
42	The estimated harvest of Kaminuriak caribou by hunters from the seven Keewatin communities, for the period October 1981 to March 1986 . . .	68
43	The estimated harvest of Beverly caribou by Baker Lake and Repulse Bay hunters, for the period October 1981 to March 1986	69
44	The estimated harvest of Wager Bay caribou by Baker Lake, Coral Harbour, Eskimo Point, Repulse Bay and Whale Cove hunters, for the period October 1981 to March 1986	70
45	The estimated harvest of polar bears by hunters from the seven Keewatin communities for the period October 1981 to March 1986	71
46	The estimated harvest of Arctic fox by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	72
47	The estimated harvest of wolves by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	73
48	The estimated harvest of ringed seals by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	74
49	The estimated harvest of bearded seals by hunters from Chesterfield Inlet, Coral Harbour, Eskimo Point, Rankin Inlet, Repulse Bay and Whale Cove, for the period October 1981 to March 1986	75
50	The estimated harvest of beluga whales by hunters from Chesterfield Inlet, Coral Harbour, Eskimo Point, Rankin Inlet, Repulse Bay and Whale Cove, for the period October 1981 to March 1986	76
51	The estimated harvest of Canada geese by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	77
52	The estimated harvest of Snow geese by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	78
53	The estimated harvest of charr by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	79
54	The estimated harvest of lake trout by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	80

LIST OF FIGURES

Figure		Page
1	Map of Keewatin District showing the seven communities surveyed during the harvest study and the zonal grid used to locate kills	81

LIST OF APPENDICES

Appendix		Page
1	Monthly percent of hunters interviewed for the seven Keewatin communities for the period October 1981 to March 1986	82
2	Monthly theoretical kill factors for the seven Keewatin communities for the period October 1981 to March 1986	83
3	Members of the Steering Committee for the period of the Keewatin Wildlife Federation Harvest Study, 1981 to 1987	84
4	Footnotes for Tables 35 to 41 . . .	85

ABSTRACT

Gamble, R.L. 1988. Native harvest of wildlife in the Keewatin Region, Northwest Territories for the period October 1985 to March 1986 and a summary for the entire period of the harvest study from October 1981 to March 1986. Can. Data Rep. Fish. Aquat. Sci. 688: v + 85 p.

Harvest data have been collected from Inuit hunters of the Keewatin Region from September 1981 until March 1986. The project has been run by an Inuit organization, the Keewatin Wildlife Federation, supported by funding provided through interested federal and territorial government departments. Part I (Tables 1 to 34) of this report covers the period October 1985 to March 1986 and is an update and supplement to the previous reports (Technical Reports 1282, 1543 and 1544) which cover the earlier years of the survey. Results in Part I are presented in the same format as the previous noted. Data were aggregated at a community level and fieldworkers continued to maintain a high level of performance as measured by participation of hunters in the study and a subjective judgement of the quality of data based on experience. Part II of this report (Tables 35 to 54) is a summary of the entire period of the harvest study. Tables 35 to 41 give the estimated harvest of species by community for the years of survey and the remaining tables give the harvest of selected species aggregated at a regional level.

Key words: resource management; catch statistics; domestic harvest; monitoring; food resources; country foods; terrestrial mammals; marine mammals; birds; fish; computerized harvest study; Inuit organization.

RÉSUMÉ

Gamble, R.L. 1988. Native harvest of wildlife in the Keewatin Region, Northwest Territories for the period October 1985 to March 1986 and a summary for the entire period of the harvest study from October 1981 to March 1986. Can. Data Rep. Fish. Aquat. Sci. 688: v + 85 p.

Des données sur les prises/captures ont été recueillies auprès de chasseurs inuit de la région du Keewatin pour la période de septembre 1981 à mars 1986, dans le cadre d'un programme de collecte dont un organisme inuit, la Keewatin Wildlife Federation, assure l'application. Le financement pour le projet vient des ministères fédéral et territorial en cause. La Partie I (tableaux 1 à 34) du rapport porte sur la période d'octobre 1985 à mars 1986 et constitue une mise à jour et un complément aux rapports précédents du même auteur (rapports techniques 1282, 1543 et 1544) sur les premières années de l'étude. Les résultats dans cette Partie sont présentés comme dans les rapports précédents. Les données ont été groupées par collectivité et le travail de collecte a été fait de façon excellente, comme l'indiquent la participation des chasseurs à l'étude et l'évaluation subjective des données, fondée sur l'expérience. La Partie II du rapport (tableaux 35 à 54) constitue un résumé de toute la période visée par l'étude. Les tableaux 35 à 41 présentent les prises/captures estimatives des espèces par collectivité pour la durée de l'étude tandis que les tableaux 42 à 54 renferment les chiffres des prises/captures de certaines espèces regroupés par région.

Mots-clés: gestion des ressources; statistiques des prises; chasse/pêche de subsistance; contrôle; ressources alimentaires; ressources alimentaires indigènes; mammifères terrestres; mammifères marins; oiseaux; poisson; étude des prises/captures par ordinateur; organisme inuit.

INTRODUCTION

The collection of data for the "Keewatin Region Native Harvest of Wildlife Study" began in September, 1981. Previous results have been published for the following periods: October 1981 to September 1983 (Gamble 1984); October 1983 to September 1984 (Gamble 1987a); and October 1984 to September 1985 (Gamble 1987b). Part I (Tables 1 to 34) of this report supplements and updates these previous reports for the survey period October 1985 to March 1986. Part II of this report is a summary of the entire period of the harvest study from September 1981 to March 1986. In Part II, Tables 35 to 41 give the estimated harvest of species by community for the years of survey, and Tables 42 to 54 give the harvest of selected species aggregated at a regional level. Appendix 4 gives footnotes pertinent to the tables in Part II.

Throughout this report hunter, harvester, trapper and fisherman are used as synonyms. Hunter is defined in the MATERIALS AND METHODS section below.

The main objectives of the study were to:

1) determine by survey techniques the hunter kill (i.e. harvest) by Inuit living in District of Keewatin communities and outpost camps;

2) develop an approach for the collection of timely, statistically reliable data on wildlife harvesting which could be undertaken by an agency such as the Keewatin Wildlife Federation (KWF) upon completion of the preliminary study;

3) determine the number of Inuit directly participating in subsistence harvesting in each community and to compare the proportion of harvest taken by hunters of different ages;

4) provide an estimate of the harvest sufficient to determine a measure of its value to each community as food or income, and

5) analyze and publish the data collected in a timely report and scientifically acceptable format.

The study area (Fig. 1) remained the same for the entire period of the study as reported in Gamble (1984; 1987a and b) and includes the entire Keewatin district of the Northwest Territories (approximately 386,000 km^2). This region contains seven permanent communities. Listed alphabetically (the convention followed throughout this and previous reports) they are Baker Lake, Chesterfield Inlet, Coral Harbour, Eskimo Point, Rankin Inlet, Repulse Bay and Whale Cove. Current information about these communities including population can be obtained from the NWT Data Book (1986).

MATERIALS AND METHODS

GENERAL

For the entire period of survey fieldworkers tried to include 100% of the region's hunters in their monthly data collection. The study design remained the same as originally described in Gamble (1984).

For the purpose of the study the term hunter includes all Inuit males and females over the age of 16 who hunt (they may or may not have a NWT general hunting licence), Inuit youths under 16 who hunt regularly, and some long-term residents in the area of other ethnic origin who hunt. This latter group makes up less than 1% of the total hunters in the region and also accounts for less than 1% of the animals harvested.

Harvest data were aggregated at the community level. Separate coverage of outpost camps was not necessary because Inuit hunting from such locations visited their home communities frequently during the survey period and it was possible to include their harvest together with that of community based hunters on a consistent basis.

HUMAN RESOURCES AND MATERIALS

Fieldworkers were hired in each of the seven communities to interview hunters and collect data. Duties included explaining the project to hunters; distributing the study materials (calendars and field notebooks) to hunters; keeping an up to date list of hunters; interviewing hunters beginning on the first day of each month to collect harvest statistics for the previous month and recording this information on the appropriate data sheets; making sure the data collected were as accurate as possible; and promptly forwarding a monthly report following an interview period to the Project Biologist located at Rankin Inlet.

Over the period of the survey slight changes were made to data sheets, calendars and field diaries that were distributed to fieldworkers and hunters. The project office organization was also modified. These modifications are described in Gamble 1987a and 1987b.

During the period October 1985 to March 1986 the Project Office organization remained the same as described by Gamble (1987a) and no changes were made to the data sheets, calendars and field diaries that were distributed to fieldworkers and hunters as described in Gamble (1987b).

DATA COLLECTION AND ANALYSIS

The system used to analyze harvest data and to arrive at estimates of the total hunter kill by community required several steps and remained the same as developed during the 1981-1983 preliminary study (Gamble 1984).

Beginning on the first day of each month fieldworkers began interviews so that they could divide the hunter population for each community into the survey categories defined below and list the number of animals killed per species for successful hunters who were interviewed. The monthly interval was defined as an interview period and covered the previous month of hunting. The fieldworker submitted this information to the Project Office where the data were summarized each month against a master list of hunters for individual communities and then entered into the computer. The numbers in some categories were subsequently adjusted the following month (i.e. the second month past the actual hunting episode) if acceptable reports were submitted by fieldworkers on hunters who had been interviewed after a particular interview period had passed. Acceptable reports were determined through a subjective judgement by the Project Biologist based on his experience and a comparison of late reports with the reports submitted on time.

Definition	Category
1) The number of hunters who report taking a harvest during an interview period (i.e. successful).	A
2) The number of hunters who report they were not successful in taking a harvest during an interview period (i.e. unsuccessful).	B
3) The number of hunters who report they did not hunt during an interview period (i.e. didn't hunt).	C
4) The number of hunters who were out hunting during the interview period but who were not interviewed (i.e. hunted but not interviewed).	D
5) The number of hunters who were out of the area of the harvest survey during the interview period for any reason (i.e. out of hunt area).	E
6) The number of hunters within the harvest study area during the interview period whose activities were unknown (i.e. activities unknown).	F

Subsequently the summarized monthly information contained in categories A through F was used to calculate ratios of participation and hunter success. Participation ratio refers to the percent of hunters in each community who were interviewed as part of the study in relation to the total number of hunters who could have hunted each month. Appendix 1 summarizes the monthly percent of hunters interviewed for the seven Keewatin communities for the period October 1981 to March 1986.

The hunter success ratio was applied to hunters in categories D and F to obtain an estimate of probable hunter success within these groups. The results for all categories were summed to get an estimate of total hunter success and to calculate the theoretical kill factor. This is the value by which the reported kill per species is multiplied to arrive at the estimated harvest. The theoretical kill factors calculated for each month for the seven Keewatin communities for the period October 1981 to March 1986 are given in Appendix 2.

For the purpose of this analysis four main assumptions were made:

1) The involvement of hunters in the harvest is the same for those whose activities are unknown as for those that are known.
2) The success ratio is the same for hunters who hunted in the unknown categories as for the known categories.
3) The probability of a kill of any individual animal is the same for all species when calculating the estimated harvest.
4) Reported kills are accurate.

DATA PROCESSING

The study used the basic computer programs described by Gamble (1984; 1987a and b) throughout the period of survey. The processing of harvest data was enhanced after the preliminary survey described by Gamble 1984 by the addition of programmes which allowed the species harvested to be reflected by a geographic zone system (Fig. 1) and a breakdown of the reported kill by species over the range of age groups for the hunters.

The hunter age classes used for the survey were 0-15, 16-30, 31-45, 46-60, 61-75 and 76-99 years. The age group 76-99 was used as a category for hunters with unknown ages because only 8 hunters of known age fell within this group.

ACKNOWLEDGMENTS

I thank the Keewatin Wildlife Federation, which was supportive of this study and provided valuable assistance at a regional and community level.

Thanks are also due to members of the steering committee who provided valuable criticism of my manuscript, in particular R. Peet and B. Hyman of DFO who assisted in the preparation and publication of this and the previous reports (Gamble 1984, 1987a and 1987b) produced for this study. I particularly thank the staff of the Keewatin Wildlife Federation "Native Harvest Study". Their continued efforts in conjunction with the various community fieldworkers allowed this study to be successfully completed.

I especially thank all the hunters who provided data on their harvests and acknowledge their cooperation and understanding in the face of repeated questions that were necessary.

REFERENCES

BELLROSE, F.C. 1976. Ducks, geese, and swans of North America. Stackpole Books. 540 p.

BERGER, T. 1977. Northern frontier - northern homeland. The report of the Mackenzie Valley Pipeline Inquiry, 2: 268 p.

BOND, W.A. 1975. Data on the biology of lake whitefish and lake trout from Kaminuriak Lake, District of Keewatin, N.W.T. Can. Fish. Mar. Serv. Data Rep. Serv. CEN/D-75-4: 28 p.

CARDER, G.W. 1983. Data from the commercial fishery for Arctic charr, Salvelinus alpinus (Linnaeus), in the Cambridge Bay and Rankin Inlet areas, Northwest Territories, 1981-82. Can. Data Rep. Fish Aquat. Sci. 391: v + 24 p.

DOME PETROLEUM LTD., ESSO RESOURCES CANADA, LTD., AND GULF CANADA RESOURCES INC. 1982. Beaufort Sea-Mackenzie Delta Environmental Impact Statement. Socio-economic effects 5(5): 42 p.

EARHART, C.M., and N.K. JOHNSON. 1970. Size dimorphism and food habits of North American owl. Condor 72: 251-264.

FALK, M.R., and D.V. GILLMAN. 1975. Data on the lake and round whitefish, lake cisco, northern pike, Arctic grayling and longnose sucker from the east arm of Great Slave Lake, N.W.T., 1971-74. Can. Fish. Mar. Serv. Data Rep. Ser. CEN/D-75-2: 95 p.

GAMBLE, R.L. 1984. A preliminary study of the native harvest of wildlife in the Keewatin Region, Northwest Territories, Canada. Can. Tech. Rep. Fish. Aquat. Sci. 1282: iv + 48 p.

GAMBLE, R.L. 1987a. Native harvest of wildlife in the Keewatin Region, Northwest Territories for the period October 1983 to September 1984. Can. Tech. Rep. Fish. Aquat. Sci. 1543: v + 82 p.

GAMBLE, R.L. 1987b. Native harvest of wildlife in the Keewatin Region, Northwest Territories for the period October 1984 to September 1985. Can. Tech. Rep. Fish. Aquat. Sci. 1549: v + 59 p.

KELEHER, J.J. 1964. Round weight conversion factors for Great Slave Lake fish. Fish. Res. Board Can. Manuscr. Rep. Ser. (Biol.) 773: 19 p.

MacDONALD, G., and R. FUDGE. 1979. Arctic Land Use Research Program 1978: A survey of the fisheries resources of the Kazan Upland (Southeastern District of Mackenzie, Southern District of Keewatin, N.W.T.). Can. Dep. Ind. N. Aff. Environ. Stud. 11: 161 p.

NATIVE HARVESTING RESEARCH COMMITTEE (NHRC). 1975. Research to establish present levels of harvesting by native peoples of northern Quebec. Phase 1 (1973-1975). Part II. Montreal. 230 p.

NATIVE HARVESTING RESEARCH COMMITTEE (NHRC). 1976a. Research to establish present levels of harvesting by native peoples of northern Quebec. Part II. A report on the harvesting by the James Bay Cree.

NATIVE HARVESTING RESEARCH COMMITTEE (NHRC). 1976b. Research to establish present levels of harvesting by native peoples of northern Quebec. Part II. A report on the harvestng by the Inuit of northern Quebec.

NWT DATA BOOK. 1986. A complete information guide to the Northwest Territories and its communities. M. Devine (ed.) Outcrop Ltd., The Northern Publishers, Yellowknife. 236 p.

RIEWE, R. 1977. Utilization of wildlife in the Jones Sound region by the Griese Fiord Inuit, p. 623-644. In L.C. Bliss (ed.) True Love Lowlands, Devon Island, Canada: a high Arctic ecosystem. University of Alberta Press, Edmonton.

SERGEANT, D.E., and P.S. BRODIE. 1969. Body size in white whales, Delphinapterus leucas. J. Fish. Res. Board Can. 26: 2561-2580.

STEVENS, W.J.D. 1965. Bionomics of the sandhill crane. Ph.D. Thesis. University of Saskatchewan, Saskatoon. 120 p.

THOMAS, V.G. 1982. Energetic reserves of Hudson Bay willow ptarmigan during the winter and spring. Can. J. Zool. 60: 1618-1623.

Table 1. The reported harvest by Baker Lake hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Kaminuriak	M	89	72	10			28	199
	F	74	56	44			48	222
	C	8	24	24			5	61
	U		3					3
	Subtotal	171	155	78			81	485
Beverly	M	86	54					140
	F	232	54					286
	C	12	18					30
	U	11						11
	Subtotal	341	126					467
Wager	M	96	14	108	149	178	150	695
	F	80	10	143	141	157	202	733
	C	30	9	36	59	81	65	280
	U	1	2			8	1	12
	Subtotal	207	35	287	349	424	418	1720
Other	F						3	3
	Subtotal						3	3
	Total	719	316	365	349	424	502	2675
Muskox							13	13
Arctic Fox			497	250	65	152	75	1039
Wolf				21	3	5	13	42
Arctic Charr		4		7				11
Lake Trout		798	520	469	261	317	250	2615
Whitefish sp.		70	65	35	14		7	191
Percent of hunters reporting		95.4	98.0	94.8	100.0	100.0	100.0	

[1]Categories are as follows: M means male, F means female, C means calf, and U means unknown.

Table 2. The reported harvest by Chesterfield Inlet hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Kaminuriak	M	8						8
	F	3						3
	U		10					10
	Subtotal	11	10					21
North of Chesterfield	M		16	15	20	31	23	105
	F		8	5	3	14	4	34
	U		9			4	8	21
	Subtotal		33	20	23	49	35	160
	Total	11	43	20	23	49	35	181
Muskox						2		2
Polar Bear		1	2					3
Arctic Fox			26				4	30
Red Fox				1				1
Wolf			2	7		3	2	14
Ringed Seal		14	10		9	6	14	53
Bearded Seal		1					3	4
Walrus						1	2	3
Eider						4		4
Arctic Charr			70				13	83
Lake Trout sp.			58					58
Arctic Grayling			14					14
Percent of hunters reporting		100.0	100.0	100.0	100.0	92.0	91.8	

[1]See Table 1.

Table 3. The reported harvest by Coral Harbour hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Southampton	M				1	1	4	6
	C				1			1
	U				1			1
	Total				3	1	4	8
Polar Bear		7	27	2	2	1	2	41
Arctic Fox			13	44	13	2	33	105
Wolf		1						1
Arctic Hare			11	9	5	8	5	38
Ringed Seal		49	68	47	81	75	73	393
Bearded Seal		1			5	5		11
Harp Seal		2	3					5
Walrus		1					2	3
Eider		11						11
Ptarmigan		28	205	282	839	284	222	1860
Arctic Charr		216	1009	116	27	36	21	1425
Arctic Cod						7		7
Percent of hunters reporting		73.3	85.7	80.0	85.0	84.8	81.0	

[1]See Table 1.

Table 4. The reported harvest by Eskimo Point hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Kaminuriak	M	112	29	26	22	55	75	319
	F	279	61	53	79	36	219	727
	C	59	11	8	8	4	35	125
	U	18	2				3	23
	Total	468	103	87	109	95	332	1194
Moose					1	1		2
Muskox						1		1
Polar Bear			7			2		9
Arctic Fox		7	211	142	35	81	33	509
Red Fox					2	5	1	8
Wolf			3	8	5	1	44	61
Arctic Hare							1	1
Otter					1			1
Arctic Ground Squirrel					1			1
Ringed Seal		15		4	1	8	11	39
Bearded Seal		2						2
Eider							2	2
Ptarmigan						51	20	71
Arctic Charr		55	98	75	33			261
Lake Trout		70	49	26	2	3	58	208
Whitefish sp.			56	42				98
Arctic Grayling		7				3		10
Percent of hunters reporting		87.5	91.9	91.8	94.1	85.8	89.8	

[1]See Table 1.

Table 5. The reported harvest by Rankin Inlet hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Kaminuriak	M	52	24	21	30	33	30	190
	F	64	92	28	49	59	32	324
	C	3	5		6	10	6	30
	U			4	3	1	1	9
	Subtotal	119	121	53	88	103	69	553
North of Chesterfield	M						27	27
	F						23	23
	C						7	7
	U						2	2
	Subtotal						59	59
	Total	119	121	53	88	103	128	612
Muskox							2	2
Polar Bear			10					10
Arctic Fox			50	109	66	20	8	253
Wolf					2	4	1	7
Arctic Hare		2			2	2		6
Ringed Seal		63	24	2	3	8	13	113
Bearded Seal		8					1	9
Harbour Seal		2						2
Harp Seal		1						1
Walrus							2	2
Eider		8						8
Ptarmigan		4						4
Searun Arctic Charr		385	586	290	297	104	99	1761
Landlocked Arctic Charr		30						30
Lake Trout				2	15			17
Percent of hunters reporting		95.0	98.1	97.5	97.9	96.8	97.2	

[1]See Table 1.

Table 6. The reported harvest by Repulse Bay hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Wager	M	38	38	31	36	75	31	249
	F	19	19	12	48	38	56	192
	C	2	3	1	2	3	10	21
	U						4	4
	Total	59	60	44	86	116	101	466
Polar Bear			7		2	4		13
Arctic Fox			2			1	5	8
Red Fox			3	6		1		10
Wolf			5	8	3	12	5	33
Wolverine				1				1
Arctic Hare							2	2
Ringed Seal		47	19	6	2	19	12	105
Bearded Seal		2		1			1	4
Walrus		3						3
Narwhal			4					4
Ptarmigan		10			2			12
Arctic Charr		20	551				63	634
Lake Trout		12		1			4	17
Percent of hunters reporting		71.4	83.3	97.7	90.3	95.2	100.0	

[1]See Table 1.

Table 7. The reported harvest by Whale Cove hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Kaminuriak	M	30	14	10	20	6	7	87
	F	32	29	33	73	43	41	251
	C			2	9	4	4	19
	U				3		1	4
	Total	62	43	45	105	53	53	361
Muskox								
Polar Bear			10				1	1
Arctic Fox								10
Red Fox				2	2			4
Arctic Hare			1	1				1
Ringed Seal		10	1	6	7	2		4
Arctic Charr		10	9	6	40	3	4	31
Lake Trout		3			10	20	3	88
						9		22
Percent of hunters reporting		100.0	100.0	100.0	98.3	100.0	100.0	

[1]See Table 1.

Table 8. The estimated harvest by Baker Lake hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Kaminuriak	M	89	72	10			28	199
	F	74	56	44			48	222
	C	8	24	24			5	61
	U		3					3
	Subtotal	171	155	78			81	485
Beverly	M	86	54					140
	F	232	54					286
	C	12	18					30
	C	11						11
	Subtotal	341	126					467
Wager	M	96	14	108	149	178	150	695
	F	80	10	143	141	157	202	733
	C	30	9	36	59	81	65	280
	U	1	2			8	1	12
	Subtotal	207	35	287	349	424	418	1720
Other	F						3	3
	Subtotal						3	3
	Total	719	316	365	349	424	502	2675
Muskox							13	13
Arctic Fox			497	250	65	152	75	1039
Wolf				21	3	5	13	42
Arctic Charr		4		7				11
Lake Trout		798	520	469	261	317	250	2615
Whitefish sp.		70	65	35	14		7	191

[1]see Table 1.

Table 9. The estimated harvest by Chesterfield Inlet hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Kaminuriak	M	8						8
	F	3						3
	U		10					10
	Subtotal	11	10					21
North of Chesterfield	M		16	15	20	31	23	105
	F		8	5	3	14	4	34
	U		9			4	8	21
	Subtotal		33	20	23	49	35	160
	Total	11	43	20	23	49	35	181
Muskox						2		2
Polar Bear		1	2					3
Arctic Fox			26				4	30
Red Fox				1				1
Wolf			2	7		3	2	14
Ringed Seal		14	10		9	6	14	53
Bearded Seal		1					3	4
Walrus						1	2	3
Eider						4		4
Arctic Charr			70				13	83
Lake Trout			58					58
Arctic Grayling			14					14

[1]See Table 1.

Table 10. The estimated harvest by Coral Harbour hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Southampton	M				1	1	4	6
	C				1			1
	U				1			1
	Total				3	1	4	8
Polar Bear		11	35	3	2	1	2	54
Arctic Fox			17	69	13	2	35	136
Wolf		2						2
Arctic Hare			14	14	5	9	5	47
Ringed Seal		79	86	74	81	83	78	481
Bearded Seal		2			5	6		13
Harp Seal		3	4					7
Walrus		2					2	4
Eider		18						18
Ptarmigan		45	260	443	839	312	238	2137
Arctic Charr		350	1281	182	27	40	23	1903
Arctic Cod						8		8

[1]See Table 1.

Table 11. The estimated harvest by Eskimo Point hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Kaminuriak	M	120	35	32	25	81	90	383
	F	298	74	66	90	54	264	846
	C	63	13	10	9	6	42	143
	U	19	2				4	25
	Total	500	124	108	124	141	400	1397
Moose					1	2		3
Muskox						2		2
Polar Bear			8			3		11
Arctic Fox		8	255	176	40	120	40	639
Red Fox					2	7	1	10
Wolf			4	10	6	2	53	75
Arctic Hare							1	1
Otter					1			1
Arctic Ground Squirrel							1	1
Ringed Seal		16	5		1	12	13	47
Bearded Seal		2						2
Eider							2	2
Ptarmigan						76	24	100
Arctic Charr		59	119	93	37			308
Lake Trout		75	59	32	2	4	70	242
Whitefish sp.			68	52				120
Arctic Grayling		8				4		12

[1]See Table 1.

Table 12. The estimated harvest by Rankin Inlet hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Kaminuriak	M	55	25	23	32	36	33	204
	F	67	97	31	52	65	35	347
	C	3	5		6	11	7	32
	U			4	3	1	1	9
	Subtotal	125	127	58	93	113	76	592
North of Chesterfield	M						29	29
	F						25	25
	C						8	8
	U						2	2
	Subtotal						64	64
	Total	125	127	58	93	113	140	656
Muskox							2	2
Polar Bear			10					10
Arctic Fox			52	119	70	22	9	272
Wolf					2	4	1	7
Arctic Hare		2			2	2		6
Ringed Seal		66	25	2	3	9	14	119
Bearded Seal		8					1	9
Harbour Seal		2						2
Harp Seal		1						1
Walrus							2	2
Eider		8						8
Ptarmigan		4						4
Searun Arctic Charr		404	615	316	315	114	107	1871
Landlocked Arctic Charr		32						32
Lake Trout				2	16			18

[1]See Table 1.

Table 13. The estimated harvest by Repulse Bay hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Wager	M	51	44	31	44	83	31	284
	F	25	22	12	59	42	56	216
	C	3	4	1	2	3	10	23
	U						4	4
	Total	79	70	44	105	128	101	527
Polar Bear			8		2	4		14
Arctic Fox			2			1	5	8
Red Fox			4	6		1		11
Wolf			6	8	4	13	5	36
Wolverine				1				1
Arctic Hare							2	2
Ringed Seal		62	22	6	3	21	12	126
Bearded Seal		3		1			1	5
Walrus		4						4
Narwhal			5					5
Ptarmigan		13			3			16
Arctic Charr		27	639				63	729
Lake Trout		16		1			4	21

[1]See Table 1.

Table 14. The estimated harvest by Whale Cove hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category[1]	1985 Oct.	1985 Nov.	1985 Dec.	1986 Jan.	1986 Feb.	1986 Mar.	Sum.
Caribou								
Kaminuriak	M	30	14	10	20	6	7	87
	F	32	29	33	73	43	41	251
	C			2	9	4	4	19
	U				3		1	4
	Total	62	43	45	105	53	53	361
Muskox							1	1
Polar Bear			10					10
Arctic Fox				2	2			4
Red Fox				1				1
Arctic Hare			1	1		2		4
Ringed Seal		10	1	6	7	3	4	31
Arctic Charr		10	9	6	40	20	3	88
Lake Trout		3			10	9		22

[1]See Table 1.

Table 15. The reported and estimated harvest for Baker Lake hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category[1]	Reported Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.	Estimated Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.
Caribou							
Kaminuriak	M	199	3	2	199	3	2
	F	222	2	1	222	2	1
	C	61	1	1	61	1	1
	U	3	3	0	3	3	0
	Subtotal	485	2	2	485	2	2
Reverly	M	140	3	1	140	3	1
	F	286	4	2	286	4	2
	C	30	2	1	30	2	1
	U	11	11	0	11	11	0
	Subtotal	467	3	2	467	3	2
Wager	M	695	2	1	695	2	1
	F	733	2	1	733	2	1
	C	280	1	1	280	1	1
	U	12	2	2	12	2	2
	Subtotal	1720	2	1	1720	2	1
Other	F	3	3	0	3	3	0
	Subtotal	3	3	0	3	3	0
	Total	2675	2	2	2675	2	2
Muskox	13	1	0	13	1	0	
Arctic Fox	1039	6	4	1039	6	4	
Wolf	42	3	2	42	3	2	
Arctic Charr	11	6	2	11	6	2	
Lake Trout	2615	37	11	2615	37	11	
Whitefish sp.	191	7	3	191	7	3	

[1]see Table 1.

[2]See also Tables 1 and 8.

Table 16. The reported and estimated harvest for Chesterfield Inlet hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category[1]	Reported Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.	Estimated Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.
Caribou							
Kaminuriak	M	8	3	1	8	3	1
	F	3	2	1	3	2	1
	U	10	5	1	10	5	1
	Subtotal	21	3	2	21	3	2
North of Chesterfield	M	105	3	1	105	3	1
	F	34	2	1	34	2	1
	U	21	4	2	21	4	2
	Subtotal	160	3	1	160	3	1
	Total	181	3	1	181	3	1
Muskox		2	1	0	2	1	0
Polar Bear		3	1	0	3	1	0
Arctic Fox		30	4	1	30	4	1
Red Fox		1	1	0	1	1	0
Wolf		14	2	1	14	2	1
Ringed Seal		53	3	2	53	3	2
Bearded Seal		4	1	0	4	1	0
Walrus		3	1	0	4	1	0
Eider		4	4	0	4	4	0
Arctic Charr		83	21	15	83	21	15
Lake Trout		58	19	11	58	19	11
Arctic Grayling		14	7	3	14	7	3

[1]see Table 1.

[2]See also Tables 2 and 9.

Table 17. The reported and estimated harvest for Coral Harbour hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category[1]	Reported Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.	Estimated Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.
Caribou							
Southampton	M	6	1	0	6	1	0
	C	1	1	0	1	1	0
	U	1	1	0	1	1	0
	Total	8	1	0	8	1	0
Polar Bear		41	1	0	55	1	0
Arctic Fox		105	5	6	136	6	8
Wolf		1	1	0	2	2	0
Arctic Hare		38	2	2	47	3	3
Ringed Seal		393	3	3	481	4	4
Bearded Seal		11	1	0	12	1	0
Harp Seal		5	2	1	7	2	1
Walrus		3	1	0	4	1	0
Eider		11	6	5	18	9	7
Ptarmigan		1860	17	16	2137	19	19
Arctic Charr		1425	41	37	1903	54	47
Arctic Cod		7	7	0	8	8	0

[1]see Table 1.

[2]See also Tables 3 and 10.

Table 18. The reported and estimated harvest for Eskimo Point hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category[1]	Reported Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.	Estimated Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.
Caribou							
Kaminuriak	M	319	3	2	383	3	2
	F	727	3	3	844	4	3
	C	125	2	2	143	2	2
	U	23	5	5	25	5	5
	Total	1194	3	3	1395	4	3
Moose		2	1	0	3	1	0
Muskox		1	1	0	2	2	0
Polar Bear		9	1	0	11	1	0
Arctic Fox		509	8	10	638	11	13
Red Fox		8	2	1	11	3	1
Wolf		61	4	4	74	5	5
Arctic Hare		1	1	0	1	1	0
Otter		1	1	0	1	1	0
Arctic Ground Squirrel		1	1	0	1	1	0
Ringed Seal		39	2	1	47	2	1
Bearded Seal		2	1	0	2	1	0
Eider		2	2	0	2	2	0
Ptarmigan		71	24	20	100	33	30
Arctic Charr		261	11	9	308	13	11
Lake Trout		208	8	10	243	9	11
Whitefish sp.		98	33	16	120	40	20
Arctic Grayling		10	5	2	12	6	2

[1]see Table 1.

[2]See also Tables 4 and 11.

Table 19. The reported and estimated harvest for Rankin Inlet hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category[1]	Reported Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.	Estimated Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.
Caribou							
Kaminuriak	M	190	3	2	203	3	2
	F	324	3	2	346	3	2
	C	30	2	1	32	2	1
	U	9	2	1	10	2	1
	Subtotal	553	3	2	591	3	2
North of Chesterfield	M	27	2	1	29	3	2
	F	23	2	1	25	3	1
	C	7	2	0	8	3	1
	U	2	2	0	2	2	0
	Subtotal	59	2	1	64	3	1
	Total	612	3	2	655	3	2
Muskox		2	1	0	2	1	0
Polar Bear		10	1	0	10	1	0
Arctic Fox		253	9	12	272	10	13
Wolf		7	2	1	8	2	1
Arctic Hare		6	2	1	6	2	0
Ringed Seal		113	3	3	120	3	3
Bearded Seal		9	2	1	9	2	1
Harbour Seal		2	2	0	2	2	0
Harp Seal		1	1	0	1	1	0
Walrus		2	2	0	2	2	0
Eider		8	4	3	8	4	3
Ptarmigan		4	4	0	4	4	0
Searun Arctic Charr		1761	35	26	1872	37	27
Landlocked Arctic Charr		30	30	0	32	32	0
Lake Trout		17	4	2	18	5	2

[1]see Table 1.

[2]See also Tables 5 and 12.

Table 20. The reported and estimated harvest for Repulse Bay hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category[1]	Reported Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.	Estimated Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.
Caribou							
Wager	M	249	2	1	284	3	2
	F	192	2	1	217	3	2
	C	21	2	1	23	2	1
	U	4	2	1	4	2	1
	Total	466	2	1	528	3	2
Polar Bear		13	1	0	14	1	0
Arctic Fox		8	2	1	8	2	1
Red Fox		10	1	1	11	2	1
Wolf		33	2	1	36	2	1
Wolverine		1	1	0	1	1	0
Arctic Hare		2	2	0	2	2	0
Ringed Seal		105	3	3	126	3	3
Bearded Seal		4	1	0	5	1	0
Walrus		3	2	1	4	2	1
Narwhal		4	1	0	5	1	0
Ptarmigan		12	6	4	16	8	5
Arctic Charr		634	70	42	729	81	49
Lake Trout		17	3	3	21	4	4

[1]see Table 1.

[2]See also Tables 6 and 13.

Table 21. The reported and estimated harvest for Whale Cove hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category[1]	Reported Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.	Estimated Harvest[2] Oct. 1985 - March 1986 Total	Mean	S.D.
Caribou							
Kaminuriak	M	87	2	1	87	2	1
	F	251	2	2	251	2	2
	C	19	1	0	19	1	0
	U	4	1	1	4	1	1
	Total	361	2	1	361	2	1
Muskox		1	1	0	1	1	0
Polar Bear		10	1	0	10	1	0
Arctic Fox		4	1	1	4	1	1
Red Fox		1	1	0	1	1	0
Arctic Hare		4	1	1	4	1	1
Ringed Seal		31	1	0	31	1	0
Arctic Charr		88	13	12	88	13	12
Lake Trout		22	6	3	22	6	3

[1]see Table 1.

[2]See also Tables 7 and 14.

Table 22. Monthly theoretical kill factors for seven Keewatin communities.

	1985 Oct.	Nov.	Dec.	1986 Jan.	Feb.	Mar.
Baker Lake	1.00[1]	1.00	1.00	1.00	1.00	1.00
Chesterfield Inlet	1.00	1.00	1.00	1.00	1.00	1.00
Coral Harbour	1.62	1.27	1.57	1.00	1.10	1.07
Eskimo Point	1.07	1.21	1.24	1.13	1.48	1.21
Rankin Inlet	1.05	1.05	1.09	1.06	1.10	1.08
Repulse Bay	1.33	1.16	1.00	1.23	1.10	1.00
Whale Cove	1.00	1.00	1.00	1.00	1.00	1.00

[1]Theoretical kill factors derived using the method of Gamble (1984).

Table 23. The harvest by species over the range of age for Baker Lake hunters covering the period October 1985 to March 1986.

Species	Category[1]	Number of Animals Harvested Per Age Class of Hunter 1	2	3	4	5[2]
Caribou						
Kaminuriak	M		44	69	75	11
	F	6	47	93	64	12
	C		24	14	10	8
	U			3		
	Subtotal	11	115	179	149	31
Beverly	M	2	22	56	52	8
	F	7	40	118	94	27
	C		10	7	10	3
	U				11	
	Subtotal	9	72	181	167	38
Wager	M	14	251	262	131	37
	F	10	204	274	180	65
	C	2	123	95	40	20
	U			4	8	
	Subtotal	26	578	635	359	122
Other	F		3			
	Subtotal		3			
	Total	46	786	995		
Muskox			2	5	6	
Arctic Fox		13	135	349	420	122
Wolf			5	11	26	
Arctic Charr					11	
Lake Trout			348	752	856	659
Whitefish sp.			6	58	92	35

[1]see Table 1.

[2]Age classes are as follows: 1 = 0-15
2 = 16-30
3 = 31-45
4 = 46-60
5 = 61-75

Table 24. The harvest by species over the range of age for Chesterfield Inlet hunters covering the period October 1985 to March 1986.

Species	Category[1]	Number of Animals Harvested Per Age Class of Hunter				
		1	2	3	4	5[2]
Caribou						
Kaminuriak	M		2	4	2	
	F			1	2	
	U			4	6	
	Subtotal		2	9	10	
North of Chesterfield	M		17	50	30	8
	F		1	10	23	
	U		2	10	9	
	Subtotal		20	70	62	8
	Total		22	79	72	8
Muskox			1		1	
Polar Bear			2		1	
Arctic Fox		3	12		15	
Red Fox				1		
Wolf			1	7	6	
Ringed Seal			6	14	33	
Bearded Seal				1	3	
Walrus			1		2	
Eider				4		
Arctic Charr			7	30	46	
Lake Trout				20	38	
Arctic Grayling				10	4	

[1]see Table 1.

[2]For age classes see Table 23.

Table 25. The harvest by species over the range of age for Coral Harbour hunters covering the period October 1985 to March 1986.

Species	Category[1]	Number of Animals Harvested Per Age Class of Hunter				
		1	2	3	4	5[2]
Caribou						
Southampton	M			2	4	
	C		1			
	U			1		
	Total		1	3	4	
Polar Bear		1	14	18	6	2
Arctic Fox			25	33	44	3
Wolf				1		
Arctic Hare			20	11	6	1
Ringed Seal			75	173	129	16
Bearded Seal				5	6	
Harp Seal			1	4		
Walrus			2	1		
Eider				10	1	
Ptarmigan		2	442	779	580	37
Arctic Charr			459	620	324	22
Arctic Cod				7		

[1]see Table 1.

[2]For age classes see Table 23.

Table 26. The harvest by species over the range of age for Eskimo Point hunters covering the period October 1985 to March 1986.

Species	Category[1]	Number of Animals Harvested Per Age Class of Hunter 1	2	3	4	5[2]
Caribou						
Kaminuriak	M		96	110	108	5
	F		127	373	223	4
	C		32	62	27	4
	U		5	18		
	Total		260	563	358	13
Moose				1	1	
Muskox					1	
Polar Bear			3	3	2	1
Arctic Fox			93	220	196	
Red Fox			1	4	3	
Wolf			19	40	2	
Arctic Hare			1			
Otter					1	
Arctic Ground Squirrel			1			
Ringed Seal			10	22	7	
Bearded Seal			1	1		
Eider			2			
Ptarmigan				70	1	
Arctic Charr			59	135	67	
Lake Trout			85	54	69	
Whitefish sp.					98	
Arctic Grayling			3	7		

[1]see Table 1.

[2]For age classes see Table 23.

Table 27. The harvest by species over the range of age for Rankin Inlet hunters covering the period October 1985 to March 1986.

Species	Category[1]	Number of Animals Harvested Per Age Class of Hunter 1	2	3	4	5[2]
Caribou						
Kaminuriak	M		52	61	72	5
	F		60	100	139	25
	C		4	13	13	
	U			3	5	1
	Subtotal		116	177	229	31
North of Chesterfield	M		3	16	8	
	F		2	11	10	
	C			5	2	
	U				2	
	Subtotal		5	32	22	
	Total		121	209	251	31
Muskox				2		
Polar Bear			6	3	1	
Arctic Fox			4	40	139	70
Wolf			1	6		
Arctic Hare			2	4		
Ringed Seal			23	68	18	4
Bearded Seal			3	5	1	
Harbour Seal			2			
Harp Seal					1	
Walrus				2		
Eider				7	1	
Ptarmigan					4	
Searun Arctic Charr			140	473	971	177
Landlocked Arctic Charr					30	
Lake Trout			6		9	2

[1]see Table 1.

[2]For age classes see Table 23.

Table 28. The harvest by species over the range of age for Repulse Bay hunters covering the period October 1985 to March 1986.

Species	Category[1]	Number of Animals Harvested Per Age Class of Hunter				
		1	2	3	4	5[2]
Caribou						
Wager	M		73	78	68	30
	F		72	88	19	13
	C		12	1	8	
	U		1	3		
	Total		158	170	95	43
Polar Bear			2	5	4	2
Arctic Fox			1	4	1	2
Red Fox			4		3	3
Wolf			12	18	2	1
Wolverine					1	
Arctic Hare						2
Ringed Seal			13	45	29	18
Bearded Seal				2	1	1
Walrus			1			2
Narwhal			2	2		
Ptarmigan						12
Arctic Charr			63	487	64	20
Lake Trout			5	9	1	2

[1]See Table 1.

[2]For age classes see Table 23.

Table 29. The harvest by species over the range of age for Whale Cove hunters covering the period October 1985 to March 1986.

Species	Category[1]	Number of Animals Harvested Per Age Class of Hunter				
		1	2	3	4	5[2]
Caribou						
Kaminuriak	M		23	38	21	5
	F		80	64	95	12
	C		4	1	13	1
	U		4			
	Total		111	103	129	18
Muskox			1			
Polar Bear			4	4	2	
Arctic Fox			2	1	1	
Red Fox				1		
Arctic Hare			3			1
Ringed Seal			13	8	5	5
Arctic Charr			12	55		21
Lake Trout				16		6

[1]See Table 1.

[2]For age classes see Table 23.

Table 30. Data on the distribution of hunters that were successful in obtaining a harvest expressed as a percentage over the range of age of hunters for the period October 1985 to March 1986.

Community	Range of Ages	Distribution of Successful Hunters by Month (%) Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Total by Harvest Year
Baker Lake	0-15	3.9	1.0	2.1	1.8	2.2	2.8	1.8
	16-30	33.7	37.5	34.2	37.9	38.7	40.1	43.4
	31-45	32.6	30.7	31.6	30.2	33.9	31.1	27.6
	46-60	21.5	22.4	22.1	21.3	19.4	17.9	18.4
	61-75	8.3	8.3	10.0	8.9	5.9	8.0	8.8
Number of successful hunters		181	192	190	169	186	212	272
Chesterfield Inlet	0-15	0.0	5.9	0.0	0.0	0.0	0.0	2.9
	16-30	37.5	41.2	14.3	0.0	27.8	31.3	42.9
	31-45	37.5	23.5	42.9	42.9	33.3	37.5	28.6
	46-60	25.0	23.5	28.6	42.9	38.9	31.3	22.9
	61-75	0.0	5.9	14.3	14.3	0.0	0.0	2.9
Number of successful hunters		8	17	7	7	18	16	35
Coral Harbour	0-15	0.0	2.1	0.0	0.0	0.0	0.0	1.1
	16-30	48.1	29.2	16.0	31.6	21.2	26.9	40.2
	31-45	29.6	43.8	40.0	38.6	48.5	50.0	34.8
	46-60	18.5	18.8	32.0	21.1	27.3	19.2	15.2
	61-75	3.7	6.3	12.0	8.8	3.0	3.8	8.7
Number of successful hunters		27	48	25	57	33	26	92
Eskimo Point	0-15	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	16-30	21.4	31.1	23.7	18.4	29.3	35.4	30.6
	31-45	47.6	28.9	39.5	47.4	39.0	43.9	40.8
	46-60	28.6	35.6	36.8	34.2	29.3	19.5	25.9
	61-75	2.4	4.4	0.0	0.0	2.4	1.2	2.7
Number of successful hunters		84	45	38	38	41	82	147
Rankin Inlet	0-15	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	16-30	28.2	30.6	9.5	14.3	26.7	19.4	28.6
	31-45	35.9	30.6	33.3	25.0	40.0	41.7	35.7
	46-60	30.8	36.1	42.9	53.6	30.0	33.3	28.6
	61-75	5.1	2.8	14.3	7.1	3.3	5.6	7.1
Number of successful hunters		39	36	21	28	30	36	84
Repulse Bay	0-15	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	16-30	25.0	35.3	37.0	30.0	36.7	48.7	44.1
	31-45	29.2	29.4	33.3	46.7	36.7	33.3	30.9
	46-60	25.0	20.6	18.5	16.7	23.3	12.8	14.7
	61-75	20.8	14.7	11.1	6.7	3.3	5.1	10.3
Number of successful hunters		24	34	27	30	30	39	68
Whale Cove	0-15	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	16-30	43.3	46.4	34.8	51.9	40.9	50.0	47.8
	31-45	20.0	28.6	26.1	25.9	27.3	20.0	21.7
	46-60	26.7	21.4	26.1	18.5	27.3	30.0	19.6
	61-75	10.0	3.6	13.0	3.7	4.5	0.0	10.9
Number of successful hunters		30	28	23	27	22	20	46
Regional Total	0-15	1.8	1.0	1.2	.8	1.1	1.4	.9
	16-30	31.8	35.8	29.9	32.6	34.4	37.6	39.1
	31-45	34.9	31.5	33.2	34.3	36.1	35.5	32.0
	46-60	24.4	24.5	26.0	25.0	23.9	20.2	20.6
	61-75	7.1	7.3	9.7	7.3	4.4	5.3	7.4
Number of successful hunters		393	400	331	356	360	431	744

Table 31. Edible weight values in kilograms for harvested species as calculated from various sources.

Species	Estimated Individual Weight (kg)	Reference[1]
Caribou	48.0	Berger 1977
Moose	199.0	Berger 1977
Muskox	110.0	Riewe 1977
Polar bear	158.8	Native Harvesting Research Committee 1975, 1976a or b
Black bear	45.4	Dome et al. 1982
Grizzly bear	45.4	"
Arctic hare	2.3	Native Harvesting Research Committee 1975, 1976a or b
Ringed seal	14.3	"
Bearded seal	98.4	"
Harbour seal	27.7	"
Harp seal	43.1	"
Walrus	185.1	"
Beluga[2]	(M)555.0(F)407.9	Sergeant and Brodie 1969
Narwhal[2]	(M)595.2(F)397.0	Hay (personal communication, DFO, St. Johns's, NF); Sergeant and Brodie 1969
Canada geese (Hutchinsii)	2.4	Bellrose 1976
Snow geese (Lesser)	1.6	"
Ross's geese	1.0	"
Eider (Hudson Bay)	1.5	"
Old squaw	0.5	"
Mallard	0.7	"
Ptarmigan	0.4	Thomas 1982
Sandhill crane	4.1	Stevens 1965
Snowy owl	1.8	Earhart and Johnson 1970
Swan	6.8	Bellrose 1976
Arctic charr	2.5	Carder 1983
Lake trout	2.4	Bond 1975; Keleher 1964
Whitefish sp.	2.8	"
Northern pike	2.1	MacDonald and Fudge 1979; Keleher 1964
Arctic grayling	0.9	Falk and Gillman 1975; Keleher 1964

[1]These references are listed in detail in the reference section of the report.

[2]"M" means male, "F" means female.

Table 32. Reported and estimated edible weight values (kg) for harvested species for the period October, 1985 to March, 1986. The reported and estimated edible weight values for Baker Lake, Chesterfield Inlet and Whale Cove are the same as all six theoretical kill values equalled one.

Community and Species	1985-86 Reported Harvest (kg) Total	1985-86 Estimated Harvest (kg) Total
Baker Lake		
Caribou	128400	128400
Muskox	1430	1430
Arctic Charr	28	28
Lake Trout	6276	6276
Whitefish sp.	535	535
Total	136669	136669
Chesterfield Inlet		
Caribou	8688	8688
Muskox	220	220
Polar Bear	476	476
Ringed Seal	758	758
Bearded Seal	394	394
Walrus	555	555
Eider	6	6
Arctic Charr	208	208
Lake Trout	139	139
Arctic Grayling	13	13
Total	11457	11457
Coral Harbour		
Caribou	384	402
Polar Bear	6511	8577
Arctic Hare	87	109
Ringed Seal	5620	6880
Bearded Seal	1082	1193
Harp Seal	216	304
Walrus	555	696
Eider	17	27
Ptarmigan	744	855
Arctic Charr	3563	4756
Total	18779	23799
Eskimo Point		
Caribou	57312	67141
Moose	398	519
Muskox	110	163
Polar Bear	1429	1815
Arctic Hare	2	3
Ringed Seal	558	676
Bearded Seal	197	211
Eider	3	4
Ptarmigan	28	40
Arctic Charr	653	769
Lake Trout	499	584
Whitefish sp.	274	336
Arctic Grayling	9	11
Total	61472	72272

Table 32 Cont'd.

Community and Species	1985-86 Reported Harvest (kg) Total	1985-86 Estimated Harvest (kg) Total
Rankin Inlet		
Caribou	29376	31420
Muskox	220	238
Polar Bear	1588	1667
Arctic Hare	14	15
Ringed Seal	1616	1710
Bearded Seal	886	933
Harbour Seal	55	58
Harp Seal	43	45
Walrus	370	400
Eider	12	13
Ptarmigan	2	2
Arctic Charr	4403	4680
Lake Trout	41	43
Total	38626	41224
Repulse Bay		
Caribou	22368	25270
Polar Bear	1747	2020
Arctic Hare	5	5
Ringed Seal	1502	1801
Bearded Seal	394	459
Walrus	555	739
Narwhal	1984	2302
Ptarmigan	5	6
Arctic Charr	1585	1822
Lake Trout	41	50
Total	30186	34474
Whale Cove		
Caribou	17328	17328
Muskox	110	110
Polar Bear	1588	1588
Arctic Hare	9	9
Ringed Seal	443	443
Arctic Charr	163	163
Lake Trout	53	53
Total	19694	19694

[4]Please note that rounding has caused small discrepancies in column totals.

Table 33. Reported and estimated edible weight values (kg) for four major groups of animals harvested by Keewatin communities, October, 1985 to March, 1986.

Period	Baker Lake (reported edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish	Baker Lake (estimated edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish
1985										
Oct	36633[1]	34512 (94.2)			2121 (5.8)	36633[1]	34512 (94.2)			2121 (5.8)
Nov	16598	15168 (91.4)			1430 (8.6)	16598	15168 (91.4)			1430 (8.6)
Dec	18761	17520 (93.4)			1241 (6.6)	18761	17520 (93.4)			1241 (6.6)
1986										
Jan	17418	16752 (96.2)			666 (3.8)	17418	16752 (96.2)			666 (3.8)
Feb	21113	20352 (96.4)			761 (3.6)	21113	20352 (96.4)			761 (3.6)
Mar	26146	25526 (97.6)			620 (2.4)	26146	25526 (97.6)			620 (2.4)
Total	136669	129830 (95.0)			6839 (5.0)	136669	129830 (95.0)			6839 (5.0)

[1]The theoretical kill factor (Table 22) is the value by which the reported kill per species is multiplied to arrive at the estimated harvest. In cases where this value is one then 100% of the hunters have been interviewed and the reported and estimated harvests are equal.

Table 33 Cont'd.

Period	Chesterfield Inlet (reported edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish	Chesterfield Inlet (estimated edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish
1985										
Oct	985	687 (69.7)	299 (30.3)			985	687 (69.7)	299 (30.3)		
Nov	2852	2382 (83.5)	143 (5.0)		327 (11.5)	2852	2382 (83.5)	143 (5.0)		327 (11.5)
Dec	960	960 (100.0)				960	960 (100.0)			
1986										
Jan	1233	1104 (89.6)	129 (10.4)			1233	1104 (89.6)	129 (10.4)		
Feb	2849	2572 (90.3)	271 (9.5)	6 (.2)		2849	2572 (90.3)	271 (9.5)	6 (.2)	
Mar	2579	1680 (65.2)	866 (33.6)		33 (1.3)	2579	1680 (65.2)	866 (33.6)		33 (1.3)
Total	11458	9385 (81.9)	1708 (14.9)	6 (.1)	360 (3.1)	11458	9385 (81.9)	1708 (14.9)	6 (.1)	360 (3.1)

Table 33 Cont'd.

Period	Coral Harbour (reported edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish	Coral Harbour (estimated edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish
1985										
Oct	2750	1112 (40.4)	1070 (38.9)	28 (1.0)	540 (19.6)	4455	1801 (40.4)	1734 (38.9)	45 (1.0)	875 (19.6)
Nov	8020	4313 (53.8)	1102 (13.7)	82 (1.0)	2523 (31.5)	10184	5477 (53.8)	1399 (13.7)	104 (1.0)	3204 (31.5)
Dec	1413	338 (23.9)	672 (47.6)	113 (8.0)	290 (20.5)	12218	531 (23.9)	1055 (47.6)	177 (8.0)	455 (20.5)
1986										
Jan	2527	473 (18.7)	1650 (65.3)	336 (13.3)	68 (2.7)	2527	473 (18.7)	1650 (65.3)	336 (13.3)	68 (2.7)
Feb	1994	225 (11.3)	1565 (78.5)	114 (5.7)	90 (4.5)	2193	248 (11.3)	1721 (78.5)	125 (5.7)	99 (4.5)
Mar	2077	521 (25.1)	1414 (68.1)	89 (4.3)	53 (2.5)	2222	558 (25.1)	1513 (68.1)	95 (4.3)	56 (2.5)
Total	18781	6982 (37.2)	7473 (39.8)	762 (4.0)	3564 (19.0)	23799	9088 (38.2)	9073 (38.1)	882 (3.7)	4757 (20.0)

Table 33 Cont'd.

Period	Eskimo Point (reported edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish	Eskimo Point (estimated edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish
1985										
Oct	23187	22464 (96.9)	411 (1.8)		312 (1.3)	24811	24037 (96.9)	440 (1.8)		334 (1.3)
Nov	6575	6056 (92.1)			519 (7.9)	7956	7327 (92.1)			629 (7.9)
Dec	4601	4176 (90.8)	57 (1.2)		368 (8.0)	5705	5178 (90.8)	71 (1.2)		456 (8.0)
1986										
Jan	5532	5431 (98.2)	14 (.3))		87 (1.6)	6252	6137 (98.2)	16 (.3)		99 (1.6)
Feb	5331	5187 (97.3)	114 (2.1)	20 (.4)	10 (.2)	7890	7676 (97.3)	169 (2.1)	30 (.4)	15 (.2)
Mar	16245	15938 (98.1)	157 (1.0)	11 (.1)	139 (.9)	19656	19285 (98.1)	190 (1.0)	13 (.1)	168 (.9)
Total	61471	59252 (96.4)	753 (1.2)	31 (.1)	1435 (2.3)	72270	69640 (96.4)	886 (1.2)	43 (.1)	1701 (2.4)

Table 33 Cont'd.

Period	Rankin Inlet (reported edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish	Rankin Inlet (estimated edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish
1985										
Oct	8481	5717 (67.4)	1787 (21.1)	14 (.2)	963 (11.4)	8903	6002 (67.4)	1876 (21.1)	14 (.2)	1011 (11.4)
Nov	9204	7396 (80.4)	343 (3.7)		1465 (15.9)	9664	7766 (80.4)	360 (3.7)		1538 (15.9)
Dec	3303	2544 (77.0)	29 (.9)		730 (22.1)	3600	2773 (77.0)	31 (.9)		796 (22.1)
1986										
Jan	5051	4229 (83.7)	43 (.8)		779 (15.4)	5353	4482 (83.7)	46 (.8)		825 (15.4)
Feb	5323	4949 (93.0)	114 (2.1)		260 (4.9)	5856	5444 (93.0)	126 (2.1)		286 (4.9)
Mar	7267	6364 (87.6)	655 (9.0)		248 (3.4)	7847	6873 (87.6)	707 (9.0)		267 (3.4)
Total	38629	31199 (80.8)	2971 (7.7)	14 (.0)	4445 (11.5)	41223	33340 (80.9)	3146 (7.6)	14 (.0)	4723 (11.5)

Table 33 Cont'd.

Period	Repulse Bay (reported edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish	Repulse Bay (estimated edible wt.) Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total) Terrestrial	Marine	Fowl	Fish
1985										
Oct	5771	3767 (65.3)	1894 (32.8)	5 (.1)	105 (1.8)	4339	2832 (65.3)	1424 (32.8)	4 (.1)	79 (1.8)
Nov	8661	4446 (51.3)	2617 (30.2)		1598 (18.4)	7467	3833 (51.3)	2256 (30.2)		1378 (18.4)
Dec	2298	2112 (91.9)	184 (8.0)		2 (.1)	2298	2112 (91.9)	184 (8.0)		2 (.1)
1986										
Jan	5504	5468 (99.3)	35 (.6)	1 (.0)		4474	4446 (99.3)	27 (.6)	1 (.0)	
Feb	6948	6649 (95.7)	299 (4.3)			6316	6044 (95.7)	272 (4.3)		
Mar	5290	4853 (91.7)	270 (5.1)		167 (3.2)	5290	4853 (91.7)	270 (5.1)		167 (3.2)
Total	34472	27295 (79.2)	5299 (15.4)	6 (.0)	1872 (5.4)	30184	24120 (79.9)	4433 (14.7)	5 (.0)	1626 (5.4)

Table 33 Cont'd.

Period	Whale Cove (reported edible wt.)					Whale Cove (estimated edible wt.)				
	Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)				Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			
		Terrestrial	Marine	Fowl	Fish		Terrestrial	Marine	Fowl	Fish
1985										
Oct	3151	2976 (94.4)	143 (4.5)		32 (1.0)	3151	2976 (94.4)	143 (4.5)		32 1.0)
Nov	3691	3654 (99.0)	14 (.4)		23 (.6)	3691	3654 (99.0)	14 (.4)		23 (.6)
Dec	2263	2162 (95.5)	86 (3.8)		15 (.7)	2263	2162 (95.5)	86 (3.8)		15 (.7)
1986										
Jan	5264	5040 (95.7)	100 (1.9)		124 (2.4)	5264	5040 (95.7)	100 (1.9)		124 (2.4)
Feb	2614	2549 (97.5)	43 (1.6)		22 (.8)	2614	2549 (97.5)	43 (1.6)		22 (.8)
Mar	2711	2654 (97.9)	57 (2.1)			2711	2654 (97.9)	57 (2.1)		
Total	19694	19035 (96.7)	443 (2.3)		216 (1.1)	19694	19035 (96.7)	443 (2.3)		216 (1.1)

Table 34. Age distribution of hunters for the seven Keewatin region communities for the period October 1985 to March 1986.

Community	Percentage of Hunters Per Age Category						Total Known Hunters
	0-15	16-30	31-45	46-60	61-75	76+ [1]	
Baker Lake	1.6	41.3	27.6	20.2	6.8	2.5	366
Chesterfield Inlet	1.2	51.2	25.6	17.4	3.5	1.2	86
Coral Harbour	2.4	47.6	25.6	15.2	8.5	0.6	164
Eskimo Point	0.3	42.4	33.3	17.8	6.1	0.0	309
Rankin Inlet	0.3	40.6	34.0	17.7	6.6	0.7	288
Repulse Bay	0.8	47.5	29.5	13.1	7.4	1.6	122
Whale Cove	0.0	42.0	28.4	15.9	11.4	2.3	88
Total hunters for the Keewatin District	1.0	43.3	30.0	17.6	7.0	1.2	1423

[1]This category includes hunters of unknown ages. There are only 8 hunters of known age in this group.

Table 35. The estimated harvest by Baker Lake hunters, expressed as number of animals, for the period November, 1981 to March, 1986. Data were not available for October 1981 and August and October 1982. Results for November and December, 1981 show the actual reported harvest as data were insufficient to provide good a estimate. Therefore the actual harvest was considered to be the best estimate for these months.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
CARIBOU	KAMINURIAK (H,A)	C	1982	0	0	0	0	0	0	0	.	6	.	0	0	6	0.6	10
			1985	0	0	31	24	9	0	0	0	0	8	24	24	120	10.0	12
			1986	0	0	5	.	.	.	.	.	.	.	.	.	5	1.7	3
			N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
			SUM	0	0	36	24	9	0	0	0	6	8	24	24	.	.	.
			MEAN	0	0	7	6	2	0	0	0	2	3	5	5	.	.	.
		F	1981	.	.	.	.	.	.	.	.	.	.	114	122	236	118.0	2
			1982	353	267	147	112	14	0	2	.	143	.	195	63	1296	129.6	10
			1983	201	0	42	102	33	0	23	41	203	66	27	28	766	63.8	12
			1984	0	0	0	28	0	3	28	37	21	0	2	0	119	9.9	12
			1985	0	0	87	42	34	0	14	0	61	74	56	44	412	34.3	12
			1986	0	0	48	.	.	.	.	.	.	.	.	.	48	16.0	3
			N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
			SUM	554	267	324	284	81	3	67	78	428	140	394	257	.	.	.
			MEAN	111	53	65	71	20	1	17	26	107	47	79	51	.	.	.
		M	1981	.	.	.	.	.	.	.	.	.	.	181	144	325	162.5	2
			1982	255	161	169	129	264	10	451	.	347	.	44	47	1877	187.7	10
			1983	85	0	29	232	128	0	218	175	209	26	14	51	1167	97.3	12
			1984	0	0	0	65	0	45	275	26	10	0	0	0	421	35.1	12
			1985	0	0	104	46	44	0	64	0	104	89	72	10	533	44.4	12
			1986	0	0	28	.	.	.	.	.	.	.	.	.	28	9.3	3
			N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
			SUM	340	161	330	472	436	55	1008	201	670	115	311	252	.	.	.
			MEAN	68	32	66	118	109	14	252	67	168	38	62	50	.	.	.
		U	1982	0	34	14	0	0	0	0	.	0	.	0	0	48	4.8	10
			1984	0	0	0	0	0	3	8	0	2	0	0	0	13	1.1	12
			1985	0	0	4	0	0	0	0	0	0	0	3	0	7	0.6	12
			N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
			SUM	0	34	18	0	0	3	8	0	2	0	3	0	.	.	.
			MEAN	0	7	4	0	0	1	2	0	1	0	1	0	.	.	.

For footnotes see Appendix 4.

Table 35. (cont'd)

CARIBOU WAGER (H,A)	C	1984	0	0	0	0	0	0	0	0	0	8	58	4	70	5.8	12
		1985	0	1	0	19	12	4	2	8	35	30	9	36	156	13.0	12
		1986	59	81	65	.	.	.	.	.	.	.	.	.	205	68.3	3
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	59	82	65	19	12	4	2	8	35	38	67	40	.	.	.
		MEAN	12	16	13	5	3	1	1	3	9	13	13	8	.	.	.
	F	1983	0	0	0	0	0	100	23	57	151	20	4	31	386	32.2	12
		1984	2	0	0	0	88	17	9	242	191	240	102	2	893	74.4	12
		1985	0	9	0	33	42	57	63	227	325	80	10	143	989	82.4	12
		1986	141	157	202	.	.	.	.	.	.	.	.	.	500	166.7	3
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	143	166	202	33	130	174	95	526	667	340	116	176	.	.	.
		MEAN	29	33	40	8	33	44	24	175	167	113	23	35	.	.	.
	M	1982	0	0	0	0	0	0	0	.	6	.	0	0	6	0.6	10
		1983	0	0	0	0	0	301	169	188	329	11	4	37	1039	86.6	12
		1984	0	0	0	4	388	116	111	267	310	382	117	29	1724	143.7	12
		1985	0	5	0	45	36	70	170	325	488	96	14	108	1357	113.1	12
		1986	149	178	150	.	.	.	.	.	.	.	.	.	477	159.0	3
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	149	183	150	49	424	487	450	780	1133	489	135	174	.	.	.
		MEAN	30	37	30	12	106	122	113	260	283	163	27	35	.	.	.
	U	1983	0	0	0	0	0	0	0	3	0	0	0	0	3	0.3	12
		1984	0	0	0	0	32	21	14	2	4	1	2	0	76	6.3	12
		1985	0	0	0	0	0	3	2	0	0	1	2	0	8	0.7	12
		1986	0	8	1	.	.	.	.	.	.	.	.	.	9	3.0	3
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	0	8	1	0	32	24	16	5	4	2	4	0	.	.	.
		MEAN	0	2	0	0	8	6	4	2	1	1	1	0	.	.	.

For footnotes see Appendix 4.

Table 35. (cont'd)

CARIBOU BEVERLY (H,A)	C	1983	0	5	0	0	0	0	0	0	0	0	0	0	5	0.4	12
		1984	0	0	0	0	0	0	0	0	0	16	0	68	84	7.0	12
		1985	73	65	13	6	0	0	0	0	22	12	18	0	209	17.4	12
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	73	70	13	6	0	0	0	0	22	28	18	68	.	.	.
		MEAN	15	14	3	2	0	0	0	0	6	9	4	14	.	.	.
	F	1981	.	.	.	.	.	.	.	.	.	.	21	0	21	10.5	2
		1982	11	27	36	22	0	0	0	.	.	.	29	0	127	12.7	10
		1983	0	270	98	84	38	0	21	7	50	116	290	93	1067	88.9	12
		1984	100	247	292	234	8	0	3	0	197	212	0	117	1410	117.5	12
		1985	191	148	26	10	0	0	0	0	147	232	54	0	808	67.3	12
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	302	692	452	350	46	0	24	7	396	560	394	210	.	.	.
		MEAN	60	138	90	88	12	0	6	2	99	187	79	42	.	.	.
	M	1981	.	.	.	.	.	.	.	.	.	.	21	0	21	10.5	2
		1982	0	5	8	34	9	0	10	.	17	.	12	0	95	9.5	10
		1983	0	87	153	174	281	0	163	40	42	35	173	82	1230	102.5	12
		1984	151	337	457	424	50	0	73	0	235	330	0	146	2203	183.6	12
		1985	179	164	25	13	0	0	0	0	194	86	54	0	715	59.6	12
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	330	593	643	645	340	0	246	40	488	451	260	228	.	.	.
		MEAN	66	119	129	161	85	0	62	13	122	150	52	46	.	.	.
	U	1981	.	.	.	.	.	.	.	.	.	.	60	0	60	30.0	2
		1983	0	0	0	0	0	0	0	0	0	0	4	0	4	0.3	12
		1984	0	9	0	0	2	0	1	0	5	11	0	9	37	3.1	12
		1985	17	5	3	1	0	0	0	0	0	11	0	0	37	3.1	12
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	17	14	3	1	2	0	1	0	5	22	64	9	.	.	.
		MEAN	3	3	1	0	1	0	0	0	1	7	13	2	.	.	.

For footnotes see Appendix 4.

Table 35. (cont'd)

CARIBOU OTHER	F	1982	58	0	0	0	0	0	0	.	0	.	0	0	58	5.8	10
		1986	0	0	3	.	.	.	.	.	.	.	.	.	3	1.0	3
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	58	0	3	0	0	0	0	0	0	0	0	0	.	.	.
		MEAN	12	0	1	0	0	0	0	0	0	0	0	0	.	.	.
	M	1982	60	0	0	2	0	0	0	.	0	.	0	0	62	6.2	10
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	60	0	0	2	0	0	0	0	0	0	0	0	.	.	.
		MEAN	12	0	0	1	0	0	0	0	0	0	0	0	.	.	.
	U	1982	4	0	0	0	0	0	0	.	0	.	0	0	4	0.4	10
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	4	0	0	0	0	0	0	0	0	0	0	0	.	.	.
		MEAN	1	0	0	0	0	0	0	0	0	0	0	0	.	.	.
MUSKOX (H,A)		1982	0	0	12	0	0	0	0	.	0	.	0	0	12	1.2	10
		1983	0	0	12	0	0	0	0	0	0	0	0	0	12	1.0	12
		1984	0	0	13	0	0	0	0	0	0	0	0	0	13	1.1	12
		1985	0	0	5	0	0	0	0	0	0	0	0	0	5	0.4	12
		1986	0	0	13	.	.	.	.	.	.	.	.	.	13	4.3	3
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	0	0	55	0	0	0	0	0	0	0	0	0	.	.	.
		MEAN	0	0	11	0	0	0	0	0	0	0	0	0	.	.	.
POLAR BEAR (H,S)		1985[4]	0	0	1	0	0	0	0	0	0	0	0	0	1	0.1	12
GRIZZLY BEAR (H,S)		1984	0	0	0	0	0	0	0	0	1	0	0	0	1	0.1	12
		1985	0	0	0	0	0	0	1	0	0	0	0	0	1	0.1	12
ARCTIC FOX (H,A)		1981	.	.	.	.	.	.	.	.	.	.	13	0	13	6.5	2
		1982	0	50	71	39	0	0	0	.	0	.	417	149	726	72.6	10
		1983	30	7	0	0	0	0	0	0	0	0	17	275	329	27.4	12
		1984	124	156	180	5	0	0	0	0	0	0	526	1090	2081	173.4	12
		1985	758	585	216	113	0	0	0	0	0	0	497	250	2419	201.6	12
		1986	65	152	75	.	.	.	.	.	.	.	.	.	292	97.3	3
		N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
		SUM	977	950	542	157	0	0	0	0	0	0	1470	1764	.	.	.
		MEAN	195	190	108	39	0	0	0	0	0	0	294	353	.	.	.
RED FOX (H,S)		1984	0	0	0	0	0	0	0	0	0	0	2	2	4	0.3	12
		1985	1	0	0	0	0	0	0	0	0	0	0	0	1	0.1	12

For footnotes see Appendix 4.

Table 35. (cont'd)

WOLF (H,S)	1982	4	0	16	2	0	0	0	.	0	.	6	0	28	2.8	10
	1983	0	0	3	0	2	0	0	0	0	1	3	11	20	1.7	12
	1984	0	12	25	1	0	0	0	0	0	0	11	9	58	4.8	12
	1985	7	16	18	1	0	0	0	0	0	0	0	21	63	5.3	12
	1986	3	5	13	.	.	.	.	.	.	.	.	.	21	7.0	3
	N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
	SUM	14	33	75	4	2	0	0	0	0	1	20	41	.	.	.
	MEAN	3	7	15	1	1	0	0	0	0	0	4	8	.	.	.
RINGED SEAL (H,S)	1983	0	0	0	0	1	0	0	0	0	0	0	0	1	0.1	12
	1984	0	0	0	0	2	3	0	0	1	0	0	0	6	0.5	12
CANADA GEESE (L,S)	1984	0	0	0	0	154	142	0	0	0	0	0	0	296	24.7	12
	1985	0	0	0	0	479	144	0	0	0	0	0	0	623	51.9	12
	N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
	SUM	0	0	0	0	633	286	0	0	0	0	0	0	.	.	.
	MEAN	0	0	0	0	158	72	0	0	0	0	0	0	.	.	.
SNOW GEESE (L,S)	1984	0	0	0	0	149	201	0	0	0	0	0	0	350	29.2	12
	1985	0	0	0	0	0	30	0	0	0	0	0	0	30	2.5	12
	N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
	SUM	0	0	0	0	149	231	0	0	0	0	0	0	.	.	.
	MEAN	0	0	0	0	37	58	0	0	0	0	0	0	.	.	.
PTARMIGAN (L,A)	1981	.	.	.	.	.	.	.	.	.	.	4	0	4	2.0	2
	1984	0	0	0	0	0	0	0	0	349	403	0	0	752	62.7	12
SWAN (L,S)	1985	0	0	0	0	2	4	0	0	0	0	0	0	6	0.5	12
CANADA GOOSE EGGS	1985	0	0	0	0	151	265	0	0	0	0	0	0	416	34.7	12
GOOSE EGGS	1984	0	0	0	0	0	2722	0	0	0	0	0	0	2722	226.8	12
	1985	0	0	0	0	773	1695	0	0	0	0	0	0	2468	205.7	12
	N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
	SUM	0	0	0	0	773	4417	0	0	0	0	0	0	.	.	.
	MEAN	0	0	0	0	193	1104	0	0	0	0	0	0	.	.	.

For footnotes see Appendix 4.

Table 35. (cont'd)

ARCTIC CHARR (H,S)	1981	.	.	.	.	.	.	.	.	.	.	128	0	128	64.0	2
	1984	0	0	0	0	0	138	0	0	65	0	42	2	247	20.6	12
	1985	0	0	0	0	100	57	0	0	0	4	0	7	168	14.0	12
	N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
	SUM	0	0	0	0	100	195	0	0	65	4	170	9	.	.	.
	MEAN	0	0	0	0	25	49	0	0	16	1	34	2	.	.	.
LAKE TROUT (H,A)	1981	.	.	.	.	.	.	.	.	.	.	8744	2154	10898	5449.0	2
	1982	0	0	380	400	0	0	0	.	0	.	0	0	780	78.0	10
	1983	0	0	0	0	796	0	1886	188	0	1732	545	178	5325	443.8	12
	1984	76	157	268	241	32	175	72	87	182	196	144	192	1822	151.8	12
	1985	181	43	0	0	264	596	333	59	25	798	520	469	3288	274.0	12
	1986	261	317	250	.	.	.	.	.	.	.	.	.	828	276.0	3
	N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
	SUM	518	517	898	641	1092	771	2291	334	207	2726	9953	2993	.	.	.
	MEAN	104	103	180	160	273	193	573	111	52	909	1991	599	.	.	.
WHITEFISH (H,A)	1983	0	0	0	224	53	0	0	0	0	0	0	72	349	29.1	12
	1984	50	151	144	136	19	0	17	27	22	114	37	65	782	65.2	12
	1985	129	29	0	0	0	21	54	25	9	70	65	35	437	36.4	12
	1986	14	0	7	.	.	.	.	.	.	.	.	.	21	7.0	3
	N	5	5	5	4	4	4	4	3	4	3	5	5	.	.	.
	SUM	193	180	151	360	72	21	71	52	31	184	102	172	.	.	.
	MEAN	39	36	30	90	18	5	18	17	8	61	20	34	.	.	.
ARCTIC GRAYLING (L,A)	1984	0	0	0	0	0	0	0	0	25	0	3	0	28	2.3	12
	1985	0	0	0	0	0	56	30	0	0	0	0	0	86	7.2	12
NORTHERN PIKE (L,S)	1984	0	0	0	0	0	0	0	0	25	0	0	0	25	2.1	12
FRESHWATER FISH	1982	0	142	0	0	0	0	0	.	0	.	0	0	142	14.2	10

For footnotes see Appendix 4.

Table 36. The estimated harvest by Chesterfield Inlet hunters, expressed as number of animals for the period January, February and August, 1982 through March, 1986. The actual harvest for August, 1982 is used as it is the best estimate of the harvest for this particular month.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
CARIBOU	KAMINURIAK (H,A)[6]	F[5]	1982[1]	2	11	[3]	.	.	.	.	0[2]	0	0	15	0	28	4.0	7
			1983	11	9	0	0	0	0	0	0	0	0	0	6	26	2.2	12
			1984	17	0	1	0	6	0	0	1	0	0	0	10	35	2.9	12
			1985	0	2	0	0	0	0	0	0	0	3	0	0	5	0.4	12
			N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
			SUM	30	22	1	0	6	0	0	1	0	3	15	16	.	.	.
			MEAN	6	4	0	0	2	0	0	0	0	1	4	4	.	.	.
		M	1981	.	.	.	.	.	.	.	.	.	.	.	.	.	.	0
			1982	2	6	.	.	.	.	.	6	1	0	12	0	27	3.9	7
			1983	6	0	0	0	0	3	0	5	2	0	1	11	28	2.3	12
			1984	14	0	0	0	34	0	0	5	0	0	0	10	63	5.3	12
			1985	0	7	30	3	2	0	2	0	0	8	0	0	52	4.3	12
			N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
			SUM	22	13	30	3	36	3	2	16	3	8	13	21	.	.	.
			MEAN	4	3	8	1	12	1	1	4	1	2	3	5	.	.	.
		U	1983[4]	0	7	0	0	0	0	0	0	0	0	0	0	7	0.6	12
			1984	0	0	0	0	0	0	0	2	0	0	0	6	8	0.7	12
			1985	0	0	0	0	0	0	0	0	0	0	10	0	10	0.8	12
CARIBOU	N. CHEST.	C	1982	0	0	.	.	.	.	0	0	0	1	0	0	1	0.1	8
	(H,A)	F	1982	34	2	.	.	.	.	.	0	3	9	1	21	70	10.0	7
			1983	37	50	19	50	2	2	1	2	9	3	16	0	191	15.9	12
			1984	0	23	9	4	0	1	3	7	5	8	10	0	70	5.8	12
			1985	0	0	0	3	0	0	0	0	2	0	8	5	18	1.5	12
			1986	3	14	4	.	.	.	.	.	.	.	.	.	21	7.0	3
			N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
			SUM	74	89	32	57	2	3	4	9	19	20	35	26	.	.	.
			MEAN	15	18	8	19	1	1	1	2	5	5	9	7	.	.	.
		M	1982	18	4	.	.	.	.	.	26	35	10	4	27	124	17.7	7
			1983	31	21	31	107	26	9	56	4	10	12	3	0	310	25.8	12
			1984	0	21	31	38	23	5	14	10	35	9	4	0	190	15.8	12
			1985	3	7	9	5	13	2	8	4	15	0	16	15	97	8.1	12
			1986	20	31	23	.	.	.	.	.	.	.	.	.	74	24.7	3
			N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
			SUM	72	84	94	150	62	16	78	44	95	31	27	42	.	.	.
			MEAN	14	17	24	50	21	5	26	11	24	8	7	11	.	.	.
		U	1983	0	2	0	2	0	0	0	0	0	0	0	0	4	0.3	12
			1984	0	6	0	0	0	3	3	0	0	0	3	0	15	1.3	12
			1985	0	0	0	0	6	0	0	0	3	0	9	0	18	1.5	12
			1986	0	4	8	.	.	.	.	.	.	.	.	.	12	4.0	3
			N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
			SUM	0	12	8	2	6	3	3	0	3	0	12	0	.	.	.
			MEAN	0	2	2	1	2	1	1	0	1	0	3	0	.	.	.

For footnotes see Appendix 4.

Table 36. (cont'd)

CARIBOU OTHER	M	1984	0	0	0	0	7	0	0	0	0	0	0	0	7	0.6	12
		1985	0	0	0	0	0	0	2	0	0	0	0	0	2	0.2	12
MUSKOX (H,S)		1985	0	2	0	0	0	0	0	0	0	0	0	0	2	0.2	12
		1986	0	2	0	.	.	.	.	.	.	.	.	.	2	0.7	3
POLAR BEAR (H,S)		1982	3	0	.	.	.	.	.	0	0	0	0	1	4	0.6	7
		1983	2	0	1	7	0	0	0	0	0	0	4	0	14	1.2	12
		1984	1	0	4	0	0	0	0	0	0	0	0	0	5	0.4	12
		1985	0	1	2	0	2	0	0	0	0	1	2	0	8	0.7	12
		N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
		SUM	6	1	7	7	2	0	0	0	0	1	6	1	.	.	.
		MEAN	1	0	2	2	1	0	0	0	0	0	2	0	.	.	.
ARCTIC FOX (H,A)		1982	10	15	.	.	.	.	.	0	0	0	223	92	340	48.6	7
		1983	85	73	34	69	0	0	0	0	0	0	0	26	287	23.9	12
		1984	2	3	2	2	0	0	0	0	0	24	8	18	59	4.9	12
		1985	0	5	6	0	0	0	0	0	0	0	26	0	37	3.1	12
		1986	0	0	4	.	.	.	.	.	.	.	.	.	4	1.3	3
		N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
		SUM	97	96	46	71	0	0	0	0	0	24	257	136	.	.	.
		MEAN	19	19	12	24	0	0	0	0	0	6	64	34	.	.	.
RED FOX (H,S)		1985	0	0	0	0	0	0	0	0	0	0	0	1	1	0.1	12
WOLF (H,S)		1982	0	0	.	.	.	.	.	0	0	0	0	4	4	0.6	7
		1983	0	2	0	3	1	0	0	0	0	0	2	9	17	1.4	12
		1984	6	5	0	0	0	0	0	0	0	0	0	4	15	1.3	12
		1985	0	4	0	0	0	0	0	0	0	0	2	7	13	1.1	12
		1986	0	3	2	.	.	.	.	.	.	.	.	.	5	1.7	3
		N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
		SUM	6	14	2	3	1	0	0	0	0	0	4	24	.	.	.
		MEAN	1	3	1	1	0	0	0	0	0	0	1	6	.	.	.
RINGED SEAL (H,A)		1982	7	6	.	.	.	.	.	4	29	44	2	1	93	13.3	7
		1983	2	2	7	19	8	16	20	6	11	6	5	2	104	8.7	12
		1984	0	0	3	0	7	6	4	3	7	7	0	0	37	3.1	12
		1985	1	0	7	0	6	9	2	6	2	14	10	0	57	4.8	12
		1986	9	6	14	.	.	.	.	.	.	.	.	.	29	9.7	3
		N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
		SUM	19	14	31	19	21	31	26	19	49	71	17	3	.	.	.
		MEAN	4	3	8	6	7	10	9	5	12	18	4	1	.	.	.
BEARDED SEAL (H,S)		1982	0	0	.	.	.	.	.	0	2	0	0	0	2	0.3	7
		1983	0	0	0	0	0	0	0	0	0	3	0	0	3	0.3	12
		1984	0	0	0	0	0	0	0	0	1	0	0	0	1	0.1	12
		1985	1	0	0	1	1	1	0	0	0	1	0	0	5	0.4	12
		1986	0	0	3	.	.	.	.	.	.	.	.	.	3	1.0	3

For footnotes see Appendix 4.

Table 36. (cont'd)

WALRUS (H,S)	1983	0	0	3	7	0	0	1	0	0	0	0	0	11	0.9	12
	1984	0	0	1	4	0	2	0	0	0	0	0	0	7	0.6	12
	1985	2	4	2	0	0	8	0	0	0	0	0	0	16	1.3	12
	1986	0	1	2	.	.	.	.	.	.	.	.	.	3	1.0	3
	N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
	SUM	2	5	8	11	0	10	1	0	0	0	0	0	.	.	.
	MEAN	0	1	2	4	0	3	0	0	0	0	0	0	.	.	.
BELUGA (H,A)	1981	.	.	.	.	.	.	.	.	.	0	0	0	0	0.0	3
	1982	0	0	.	.	.	.	.	5	3	0	0	0	8	1.1	7
	1983	0	0	0	0	0	0	0	7	0	0	0	0	7	0.6	12
	1984	0	0	0	0	0	0	4	4	4	0	0	0	12	1.0	12
	1985	0	0	0	0	0	0	0	16	16	0	0	0	32	2.7	12
	N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
	SUM	0	0	0	0	0	0	4	32	23	0	0	0	.	.	.
	MEAN	0	0	0	0	0	0	1	8	6	0	0	0	.	.	.
CANADA GEESE (L,S)	1984	0	0	0	0	0	8	0	0	0	0	0	0	8	0.7	12
	1985	0	0	0	0	0	30	0	0	0	0	0	0	30	2.5	12
SNOW GEESE (L,S)	1981	.	.	.	.	.	.	.	.	.	0	0	0	0	0.0	3
	1982	0	0	.	.	.	.	.	13	7	19	0	0	39	5.6	7
	1985	0	0	0	0	0	0	0	0	9	0	0	0	9	0.8	12
EIDER (L,S)	1981	.	.	.	.	.	.	.	.	.	0	0	0	0	0.0	3
	1982	0	0	.	.	.	.	.	0	0	26	0	0	26	3.7	7
	1983	0	0	0	0	0	6	0	0	0	0	0	0	6	0.5	12
	1984	0	0	0	0	0	1	0	0	0	0	0	0	1	0.1	12
	1986	0	4	0	.	.	.	.	.	.	.	.	.	4	1.3	3
CANADA GOOSE EGGS	1984	0	0	0	0	0	0	0	2	0	0	0	0	2	0.2	12
DUCK EGGS	1984	0	0	0	0	0	8	0	0	0	0	0	0	8	0.7	12
OTHER WATERFOWL EGGS	1984	0	0	0	0	0	6	0	0	0	0	0	0	6	0.5	12

For footnotes see Appendix 4.

Table 36. (cont'd)

ARCTIC CHARR (H,A)	1982	76	0	.	.	.	.	.	0	0	0	0	0	76	10.9	7
	1983	0	0	0	0	0	55	43	55	0	0	0	0	153	12.8	12
	1984	0	0	0	0	1	0	15	414	50	0	0	0	480	40.0	12
	1985	0	0	0	0	0	8	0	0	0	0	70	0	78	6.5	12
	1986	0	0	13	.	.	.	.	.	.	.	.	.	13	4.3	3
	N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
	SUM	76	0	13	0	1	63	58	469	50	0	70	0	.	.	.
	MEAN	15	0	3	0	0	21	19	117	13	0	18	0	.	.	.
LAKE TROUT (H,A)	1982	0	19	.	.	.	.	.	202	0	26	26	0	273	39.0	7
	1983	0	0	30	131	41	72	8	0	0	0	43	0	325	27.1	12
	1984	0	0	0	1	59	26	0	0	0	111	0	0	197	16.4	12
	1985	0	0	0	2	31	22	0	0	0	0	58	0	113	9.4	12
	N	5	5	4	3	3	3	3	4	4	4	4	4	.	.	.
	SUM	0	19	30	134	131	120	8	202	0	137	127	0	.	.	.
	MEAN	0	4	8	45	44	40	3	51	0	34	32	0	.	.	.
ARCTIC GRAYLING (L,A)	1985	0	0	0	0	0	0	0	0	0	0	14	0	14	1.2	12
SCULPINS (L,A)	1984	0	0	0	0	0	1	0	0	0	0	0	0	1	0.1	12

For footnotes see Appendix 4.

Table 37. The estimated harvest by Coral Harbour hunters, expressed as number of animals, for the period February, 1982 and June 1982 through April, 1983, and October, 1983 to March, 1986. The actual harvest was used as the best estimate of the harvest for October, November, December, 1982, and October, November, December, 1983, and January, 1984.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
CARIBOU	KAMINURIAK[5]	F	1985	0[2]	0	0	6	0	0	0	0	0	0	0	0	6	0.5	12
	(H,S)[6]	M	1985	0	0	0	3	0	0	0	0	0	0	0	0	3	0.3	12
		U	1984[4]	0	0	4	0	0	0	0	0	0	0	0	0	4	0.3	12
CARIBOU	WAGER	F	1984[1]	0	0	0	78	0	0	0	3	0	0	0	0	81	6.8	12
	(H,S)		1985	0	0	0	50	63	0	0	0	0	0	0	0	113	9.4	12
			N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
			SUM	0	0	0	128	63	0	0	3	0	0	0	0	.	.	.
			MEAN	0	0	0	43	32	0	0	1	0	0	0	0	.	.	.
		M	1984	0	0	0	60	0	0	0	9	0	0	0	0	69	5.8	12
			1985	0	0	0	4	27	0	0	0	0	0	0	0	31	2.6	12
			N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
			SUM	0	0	0	64	27	0	0	9	0	0	0	0	.	.	.
			MEAN	0	0	0	21	14	0	0	2	0	0	0	0	.	.	.
		U	1984	0	0	0	55	0	0	0	0	0	0	0	0	55	4.6	12
			1985	0	0	0	30	36	0	0	0	0	0	0	0	66	5.5	12
			N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
			SUM	0	0	0	85	36	0	0	0	0	0	0	0	.	.	.
			MEAN	0	0	0	28	18	0	0	0	0	0	0	0	.	.	.

For footnotes see Appendix 4.

Table 37. (cont'd)

CARIBOU COATES (H,S)	F	1982	3 .	0	.	.	.	5	7	14	0	0	0	0	26	3.3	8
		1983	0	0	0	0	.	.	.	57	13	0	0	0	70	7.8	9
		1984	9	0	0	0	0	0	0	0	0	0	0	0	9	0.8	12
		1985	0	0	0	0	0	0	5	23	0	0	0	0	28	2.3	12
		N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
		SUM	9	0	0	0	0	5	12	94	13	0	0	0	.	.	.
		MEAN	2	0	0	0	0	2	4	24	3	0	0	0	.	.	.
	M	1982	.	0	.	.	.	0	12	11	0	0	0	0	23	2.9	8
		1983	0	0	0	0	.	.	.	70	12	0	0	0	82	9.1	9
		1984	16	0	0	0	0	0	0	0	0	0	0	0	16	1.3	12
		1985	0	0	0	0	0	0	41	65	0	0	0	0	106	8.8	12
		N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
		SUM	16	0	0	0	0	0	53	146	12	0	0	0	.	.	.
		MEAN	4	0	0	0	0	0	18	37	3	0	0	0	.	.	.
	U	1983	0	0	0	0	.	.	.	37	6	1	0	10	54	6.0	9
		1984	25	0	0	0	0	0	0	0	0	0	0	0	25	2.1	12
		1985	0	0	0	0	0	0	12	3	0	0	0	0	15	1.3	12
		N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
		SUM	25	0	0	0	0	0	12	40	6	1	0	10	.	.	.
		MEAN	6	0	0	0	0	0	4	10	2	0	0	3	.	.	.
CARIBOU SOUTHAMPTON	C	1986	1	0	0	.	.	.	.	.	.	.	.	.	1	0.3	3
(H,S)	F	1982	.	0	.	.	.	0	0	0	1	2	1	0	4	0.5	8
		1984	0	0	0	16	21	0	0	62	37	3	0	0	139	11.6	12
		1985	0	0	0	7	0	0	0	0	0	0	0	0	7	0.6	12
		N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
		SUM	0	0	0	23	21	0	0	62	38	5	1	0	.	.	.
		MEAN	0	0	0	8	11	0	0	16	10	1	0	0	.	.	.
	M	1982	.	0	.	.	.	0	10	0	10	2	2	10	34	4.3	8
		1983	0	3	0	0	.	.	.	1	1	0	4	0	9	1.0	9
		1984	0	0	0	6	23	2	3	98	39	7	3	0	181	15.1	12
		1985	0	6	1	6	15	0	0	4	2	0	0	0	34	2.8	12
		1986	1	1	4	.	.	.	.	.	.	.	.	.	6	2.0	3
		N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
		SUM	1	10	5	12	38	2	13	103	52	9	9	10	.	.	.
		MEAN	0	2	1	4	19	1	4	26	13	2	2	3	.	.	.
	U	1982	.	0	.	.	.	0	0	0	0	0	1	0	1	0.1	8
		1983	0	0	0	0	.	.	.	1	0	0	0	0	1	0.1	9
		1984	0	0	0	0	3	0	0	76	2	0	0	0	81	6.8	12
		1985	0	0	0	0	3	0	0	4	0	0	0	0	7	0.6	12
		1986	1	0	0	.	.	.	.	.	.	.	.	.	1	0.3	3
		N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
		SUM	1	0	0	0	6	0	0	81	2	0	1	0	.	.	.
		MEAN	0	0	0	0	3	0	0	20	1	0	0	0	.	.	.

For footnotes see Appendix 4.

Table 37. (cont'd)

CARIBOU OTHER	M	1985	1	0	0	0	0	0	0	0	0	0	0	0	1	0.1	12
POLAR BEAR (H,A)		1982	.	0	.	.	.	2	0	0	0	1	6	5	14	1.8	8
		1983	0	0	0	0	.	.	.	4	2	20	9	1	36	4.0	9
		1984	4	0	0	0	0	0	0	0	0	32	9	0	45	3.8	12
		1985	1	0	3	4	4	0	0	0	0	11	35	3	61	5.1	12
		1986	2	1	2	.	.	.	.	.	.	.	.	.	5	1.7	3
		N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
		SUM	7	1	5	4	4	2	0	4	2	64	59	9	.	.	.
		MEAN	2	0	1	1	2	1	0	1	1	16	15	2	.	.	.
ARCTIC FOX (H,A)		1982	.	50	.	.	.	0	0	0	0	166	167	103	486	60.8	8
		1983	117	160	224	17	.	.	.	23	21	0	126	51	739	82.1	9
		1984	85	55	89	123	0	0	0	0	0	0	109	123	584	48.7	12
		1985	29	115	76	74	12	0	0	0	0	0	17	69	392	32.7	12
		1986	13	2	35	.	.	.	.	.	.	.	.	.	50	16.7	3
		N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
		SUM	244	382	424	214	12	0	0	23	21	166	419	346	.	.	.
		MEAN	61	76	106	71	6	0	0	6	5	42	105	87	.	.	.
WOLF (H,S)		1984	0	0	0	0	1	0	0	0	0	0	0	0	1	0.1	12
		1985	0	0	0	4	4	0	0	0	0	2	0	0	10	0.8	12
ARCTIC HARE (L,S)		1982	.	0	.	.	.	0	0	0	0	0	9	9	18	2.3	8
		1983	0	0	9	0	.	.	.	0	0	0	0	0	9	1.0	9
		1984	0	0	0	0	0	0	0	0	0	0	0	3	3	0.3	12
		1985	2	0	1	7	1	0	0	0	0	0	14	14	39	3.3	12
		1986	5	9	5	.	.	.	.	.	.	.	.	.	19	6.3	3
		N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
		SUM	7	9	15	7	1	0	0	0	0	0	23	26	.	.	.
		MEAN	2	2	4	2	1	0	0	0	0	0	6	7	.	.	.

For footnotes see Appendix 4.

Table 37. (cont'd)

RINGED SEAL (H,A)	1982	.	10	.	.	.	278	107	108	52	42	30	52	679	84.9	8
	1983	67	62	40	22	.	.	.	192	49	59	87	13	591	65.7	9
	1984	134	101	55	41	57	142	74	20	45	46	20	21	756	63.0	12
	1985	89	93	19	22	30	113	83	18	33	79	86	74	739	61.6	12
	1986	81	83	78	.	.	.	.	.	.	.	.	.	242	80.7	3
	N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
	SUM	371	349	192	85	87	533	264	338	179	226	223	160	.	.	.
	MEAN	93	70	48	28	44	178	88	85	45	57	56	40	.	.	.
BEARDED SEAL (H,S)	1982	.	16	.	.	.	2	3	3	10	1	0	0	35	4.4	8
	1983	0	0	3	6	.	.	.	20	33	2	4	1	69	7.7	9
	1984	4	7	17	2	0	5	12	4	10	2	0	3	66	5.5	12
	1985	6	5	8	2	0	0	6	14	6	2	0	0	49	4.1	12
	1986	5	6	0	.	.	.	.	.	.	.	.	.	11	3.7	3
	N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
	SUM	15	34	28	10	0	7	21	41	59	7	4	4	.	.	.
	MEAN	4	7	7	3	0	2	7	10	15	2	1	1	.	.	.
HARBOUR SEAL (H,S)	1983	0	0	0	0	.	.	.	3	0	0	0	0	3	0.3	9
	1985	0	0	0	0	0	0	0	4	2	0	0	0	6	0.5	12
HARP SEAL (H,S)	1982	.	0	.	.	.	0	10	58	37	0	0	0	105	13.1	8
	1983	0	0	0	0	.	.	.	46	18	0	0	0	64	7.1	9
	1984	4	3	0	0	0	0	1	13	3	5	0	0	29	2.4	12
	1985	0	0	0	0	0	2	2	3	0	3	4	0	14	1.2	12
	N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
	SUM	4	3	0	0	0	2	13	120	58	8	4	0	.	.	.
	MEAN	1	1	0	0	0	1	4	30	15	2	1	0	.	.	.
UNKNOWN SEAL	1982	.	32	.	.	.	0	0	0	0	0	0	0	32	4.0	8
	1984	0	1	0	0	0	0	0	0	0	0	0	0	1	0.1	12
WALRUS (H,A)	1982	.	2	.	.	.	9	10	19	12	5	1	1	59	7.4	8
	1983	3	9	3	6	.	.	.	21	12	5	0	0	59	6.6	9
	1984	4	0	1	0	6	0	3	11	14	2	0	0	41	3.4	12
	1985	1	0	1	2	0	0	6	9	8	2	0	0	29	2.4	12
	1986	0	0	2	.	.	.	.	.	.	.	.	.	2	0.7	3
	N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
	SUM	8	11	7	8	6	9	19	60	46	14	1	1	.	.	.
	MEAN	2	2	2	3	3	3	6	15	12	4	0	0	.	.	.

For footnotes see Appendix 4.

Table 37. (cont'd)

BELUGA (M,A)	1982	.	0	.	.	.	0	17	75	32	0	0	1	125	15.6	8
	1983	0	0	0	0	.	.	.	77	48	2	1	0	128	14.2	9
	1984	2	1	0	0	0	3	33	50	24	8	0	0	121	10.1	12
	1985	0	0	0	0	0	0	15	60	13	0	0	0	88	7.3	12
	N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
	SUM	2	1	0	0	0	3	65	262	117	10	1	1	.	.	.
	MEAN	1	0	0	0	0	1	22	66	29	3	0	0	.	.	.
CANADA GEESE (L,S)	1982	.	0	.	.	.	41	407	199	0	0	0	9	656	82.0	8
	1983	0	0	0	0	.	.	.	9	10	0	0	0	19	2.1	9
	1984	0	0	0	0	13	101	0	23	0	0	0	0	137	11.4	12
	1985	0	0	0	0	30	25	0	0	0	0	0	0	55	4.6	12
	N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
	SUM	0	0	0	0	43	167	407	231	10	0	0	9	.	.	.
	MEAN	0	0	0	0	22	56	136	58	3	0	0	2	.	.	.
SNOW GEESE (M,A)	1982	.	0	.	.	.	4345	0	0	21	0	0	22	4388	548.5	8
	1983	0	0	0	0	.	.	.	203	12	0	0	0	215	23.9	9
	1984	3	21	0	5	322	5063	0	30	113	0	0	0	5557	463.1	12
	1985	0	0	0	0	118	3798	24	301	12	0	0	0	4253	354.4	12
	N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
	SUM	3	21	0	5	440	13206	24	534	158	0	0	22	.	.	.
	MEAN	1	4	0	2	220	4402	8	134	40	0	0	6	.	.	.
ROSS'S GEESE (L,S)	1982	.	0	.	.	.	248	7	11	1	0	0	0	267	33.4	8
BRANT GEESE (L,S)	1984	0	0	0	0	4	1	0	0	0	0	0	0	5	0.4	12
UNKNOWN GEESE	1984	0	0	0	0	3	76	0	0	0	0	0	0	79	6.6	12
	1985	0	0	0	0	0	309	0	1	0	0	0	0	310	25.8	12
OLD SQUAW (L,S)	1982	.	0	.	.	.	0	0	0	1	0	0	0	1	0.1	8
GUILLEMOT (L,S)	1982	.	0	.	.	.	0	0	0	3	0	0	0	3	0.4	8

For footnotes see Appendix 4.

Table 37. (cont'd)

EIDER	(L,A)	1982	.	0	.	.	.	217	0	28	82	0	0	0	327	40.9	8
		1983	0	0	0	0	.	.	.	31	20	12	3	0	66	7.3	9
		1984	0	0	0	0	14	6	0	0	18	11	0	0	49	4.1	12
		1985	0	0	0	0	0	0	0	0	11	18	0	0	29	2.4	12
		N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
		SUM	0	0	0	0	14	223	0	59	131	41	3	0	.	.	.
		MEAN	0	0	0	0	7	74	0	15	33	10	1	0	.	.	.
PTARMIGAN	(H,A)	1982	.	113	.	.	.	171	0	17	165	137	151	167	921	115.1	8
		1983	82	28	96	0	.	.	.	98	90	134	99	127	754	83.8	9
		1984	158	152	33	283	183	0	0	0	100	3	66	108	1086	90.5	12
		1985	33	29	175	254	206	10	0	0	17	45	260	443	1472	122.7	12
		1986	839	312	238	.	.	.	.	.	.	.	.	.	1389	463.0	3
		N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
		SUM	1112	634	542	537	389	181	0	115	372	319	576	845	.	.	.
		MEAN	278	127	136	179	195	60	0	29	93	80	144	211	.	.	.
SNOWY OWL	(L,S)	1982	.	2	.	.	.	0	0	0	0	0	0	0	2	0.3	8
UNKNOWN FOWL		1982	.	0	.	.	.	5	0	0	0	0	0	0	5	0.6	8
		1983	0	0	0	0	.	.	.	2	0	0	0	0	2	0.2	9
		1984	0	2	0	0	0	0	0	0	0	0	0	0	2	0.2	12
SWAN	(L,S)	1982	.	0	.	.	.	2	0	0	0	0	0	0	2	0.3	8
		1984	0	0	0	0	6	0	0	0	0	0	0	0	6	0.5	12
CANADA GOOSE EGGS		1984	0	0	0	0	0	71	0	0	0	0	0	0	71	5.9	12
SNOW GOOSE EGGS		1984	0	0	0	0	0	10290	0	0	0	0	0	0	10290	857.5	12
BRANT EGGS		1983	0	0	0	0	0	0	0	3	0	0	0	0	3	0.3	12
GOOSE EGGS		1984	0	0	0	0	0	30	0	0	0	0	0	0	30	2.5	12
		1985	0	0	0	0	0	3954	0	0	0	0	0	0	3954	329.5	12

For footnotes see Appendix 4.

Table 37. (cont'd)

ARCTIC CHARR (H,A)	1982	.	318	.	.	.	1504	882	1546	26	12	53	0	4341	542.6	8
	1983	0	0	0	0	.	.	.	1066	140	616	366	57	2245	249.4	9
	1984	300	12	2	210	108	370	278	690	29	1110	140	484	3733	311.1	12
	1985	0	202	331	0	138	1093	867	817	11	350	1281	182	5272	439.3	12
	1986	27	40	23	.	.	.	.	.	.	.	.	.	90	30.0	3
	N	4	5	4	3	2	3	3	4	4	4	4	4	.	.	.
	SUM	327	572	356	210	246	2967	2027	4119	206	2088	1840	723	.	.	.
	MEAN	82	114	89	70	123	989	676	1030	52	522	460	181	.	.	.
LAKE TROUT (L,S)	1982	.	0	.	.	.	0	0	0	0	0	0	9	9	1.1	8
	1983	0	0	410	0	.	.	.	0	0	0	0	0	410	45.6	9
	1985	0	0	0	0	0	7	0	0	0	0	0	0	7	0.6	12
FRESHWATER FISH	1984	0	0	0	0	0	0	0	19	0	0	0	0	19	1.6	12
ARCTIC COD (L,A)	1982	.	0	.	.	.	18	0	0	0	0	0	0	18	2.3	8
	1984	0	0	0	0	0	150	16	4	0	0	0	0	170	14.2	12
	1986	0	8	0	.	.	.	.	.	.	.	.	.	8	2.7	3
SCULPINS (L,A)	1985	0	0	0	0	0	5	0	0	0	0	0	0	5	0.4	12

For footnotes see Appendix 4.

Table 38. The estimated harvest by Eskimo Point hunters, expressed as number of animals, for the period October, 1981 to March, 1986.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	$\bar{x}$	NUM
CARIBOU	KAMINURIAK (H,A)[6]	C[5]	1981[1]	.	.[3]	.	.	.	.	.	.	.	0[2]	62	11	73	24.3	3
			1982	0	0	0	0	0	0	3	27	91	16	24	3	164	13.7	12
			1983	16	4	4	1	2	2	1	11	20	16	8	0	85	7.1	12
			1984	0	0	0	2	1	0	0	15	47	38	36	6	145	12.1	12
			1985	18	0	7	4	6	2	32	32	24	63	13	10	211	17.6	12
			1986	9	6	42	.	.	.	.	.	.	.	.	.	57	19.0	3
			N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
			SUM	43	10	53	7	9	4	36	85	182	133	143	30	.	.	.
			MEAN	9	2	11	2	2	1	9	21	46	27	29	6	.	.	.
		F	1981	.	.	.	.	.	.	.	.	.	235	350	62	647	215.7	3
			1982	58	67	136	184	28	0	35	121	226	78	57	58	1048	87.3	12
			1983	105	94	235	160	101	2	22	72	155	195	50	115	1306	108.8	12
			1984	157	292	333	255	14	4	17	22	148	65	60	55	1422	118.5	12
			1985	98	136	102	156	27	8	7	11	62	298	74	66	1045	87.1	12
			1986	90	54	264	.	.	.	.	.	.	.	.	.	408	136.0	3
			N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
			SUM	508	643	1070	755	170	14	81	226	591	871	591	356	.	.	.
			MEAN	102	129	214	189	43	4	20	57	148	174	118	71	.	.	.
		M	1981	.	.	.	.	.	.	.	.	.	196	121	80	397	132.3	3
			1982	102	131	90	84	30	10	265	302	337	173	31	19	1574	131.2	12
			1983	23	10	46	34	38	25	98	136	128	122	44	33	737	61.4	12
			1984	18	46	119	61	6	7	151	67	234	23	36	51	819	68.3	12
			1985	44	43	27	54	56	56	97	101	187	120	35	32	852	71.0	12
			1986	25	81	90	.	.	.	.	.	.	.	.	.	196	65.3	3
			N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
			SUM	212	311	372	233	130	98	611	606	886	634	267	215	.	.	.
			MEAN	42	62	74	58	33	25	153	152	222	127	53	43	.	.	.
		U	1981	.	.	.	.	.	.	.	.	.	24	6	3	33	11.0	3
			1982	25	18	32	28	12	0	11	80	76	22	9	23	336	28.0	12
			1983	45	27	25	41	5	1	23	19	10	32	10	5	243	20.3	12
			1984	18	17	13	18	12	9	5	9	31	2	2	43	179	14.9	12
			1985	6	32	7	14	34	17	14	8	0	19	2	0	153	12.8	12
			1986	0	0	4	.	.	.	.	.	.	.	.	.	4	1.3	3
			N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
			SUM	94	94	81	101	63	27	53	116	117	99	29	74	.	.	.
			MEAN	19	19	16	25	16	7	13	29	29	20	6	15	.	.	.
CARIBOU	WAGER (H,S)	M	1983[4]	0	0	0	0	0	0	0	0	1	0	0	0	1	0.1	12

For footnotes see Appendix 4.

Table 38. (cont'd)

MOOSE (H,S)	1982	0	0	0	0	1	0	0	0	0	0	0	0	1	0.1	12
	1984	0	0	3	0	2	0	0	0	0	0	0	0	5	0.4	12
	1986	1	2	0	.	.	.	.	.	.	.	.	.	3	1.0	3
MUSKOX (H,S)	1985	0	1	0	0	0	0	0	0	0	0	0	0	1	0.1	12
	1986	0	2	0	.	.	.	.	.	.	.	.	.	2	0.7	3
POLAR BEAR (H,A)	1981	.	.	.	.	.	.	.	.	.	0	10	0	10	3.3	3
	1982	0	0	0	4	0	0	0	0	0	0	13	0	17	1.4	12
	1983	0	0	1	1	0	0	0	0	0	7	12	0	21	1.8	12
	1984	0	0	1	0	1	0	0	0	0	0	13	0	15	1.3	12
	1985	0	0	0	0	0	0	0	0	0	0	8	0	8	0.7	12
	1986	0	0	3	.	.	.	.	.	.	.	.	.	3	1.0	3
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	0	0	5	5	1	0	0	0	0	7	56	0	.	.	.
	MEAN	0	0	1	1	0	0	0	0	0	1	11	0	.	.	.
ARCTIC FOX (H,A)	1981	.	.	.	.	.	.	.	.	.	0	137	90	227	75.7	3
	1982	40	58	48	172	0	0	0	0	0	24	1487	469	2298	191.5	12
	1983	210	62	80	27	6	0	0	0	0	0	40	103	528	44.0	12
	1984	117	142	163	55	1	0	0	0	0	0	227	263	968	80.7	12
	1985	28	51	262	186	0	0	0	0	0	8	255	176	966	80.5	12
	1986	40	120	40	.	.	.	.	.	.	.	.	.	200	66.7	3
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	435	433	593	440	7	0	0	0	0	32	2146	1101	.	.	.
	MEAN	87	87	119	110	2	0	0	0	0	6	429	220	.	.	.
RED FOX (H,S)	1981	.	.	.	.	.	.	.	.	.	0	0	13	13	4.3	3
	1982	0	0	2	2	0	0	0	0	0	0	28	3	35	2.9	12
	1983	5	0	3	6	0	0	0	0	0	0	2	0	16	1.3	12
	1984	1	17	9	4	0	0	0	0	0	0	7	1	39	3.3	12
	1985	0	1	0	2	0	0	0	0	0	0	0	0	3	0.3	12
	1986	2	7	1	.	.	.	.	.	.	.	.	.	10	3.3	3
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	8	25	15	14	0	0	0	0	0	0	37	17	.	.	.
	MEAN	2	5	3	4	0	0	0	0	0	0	7	3	.	.	.
WOLF (H,S)	0	0	0	0	0	0	0	0	0	0	0	0	10	10	0.8	12
	1982	0	0	2	22	0	0	0	0	0	0	1	0	25	2.1	12
	1983	0	0	0	12	8	0	0	0	0	2	4	5	31	2.6	12
	1984	0	2	12	30	2	0	0	0	0	0	3	1	50	4.2	12
	1985	0	0	3	32	2	0	0	0	0	0	4	10	51	4.3	12
	1986	6	2	53	.	.	.	.	.	.	.	.	.	61	20.3	3
	N	6	6	6	5	5	5	5	5	5	5	5	5	.	.	.
	SUM	6	4	70	96	12	0	0	0	0	2	12	16	.	.	.
	MEAN	1	1	12	19	2	0	0	0	0	0	2	3	.	.	.
MARTIN (H,S)	1982	0	0	0	0	0	0	0	0	0	0	0	1	1	0.1	12

For footnotes see Appendix 4.

Table 38. (cont'd)

OTTER (H,S)	1986	1	0	0	.	.	.	.	.	.	.	.	.	1	0.3	3
WEASEL (H,S)	1984	0	0	0	0	0	0	0	0	2	0	0	0	2	0.2	12
ARCTIC HARE (L,S)	1981	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
	1982	5	0	5	12	0	0	0	0	0	0	0	7	7	2.3	3
	1983	12	2	2	2	3	0	0	0	0	1	1	1	25	2.1	12
	1984	0	0	0	6	0	0	0	0	1	0	2	0	23	1.9	12
	1985	0	0	0	0	1	0	0	0	0	0	0	0	7	0.6	12
	1986	0	0	1	.	.	.	.	.	.	.	.	.	1	0.1	12
	N	5	5	5	4	4	4	4	4	4	5	5	5	1	0.3	3
	SUM	17	2	8	20	4	0	0	0	1	1	3	8	.	.	.
	MEAN	3	0	2	5	1	0	0	0	0	0	1	2	.	.	.
GROUND SQUIRREL (L,S)	1986	0	0	1	.	.	.	.	.	.	.	.	.	1	0.3	3
MUSKRAT (L,S)	1983	0	0	0	0	1	0	0	0	0	0	0	0	1	0.1	12
LEMMING (L,A)	1985	0	0	0	0	0	8	0	0	0	0	0	0	8	0.7	12
RINGED SEAL (H,A)	1981	.	.	.	.	.	.	.	.	.	108	2	3	113	37.7	3
	1982	1	4	3	10	5	37	20	149	70	70	4	6	379	31.6	12
	1983	3	4	0	2	2	82	31	27	12	110	36	0	309	25.8	12
	1984	17	29	20	39	36	40	44	13	132	133	0	0	503	41.9	12
	1985	0	3	0	1	35	42	10	3	7	16	5	0	122	10.2	12
	1986	1	12	13	.	.	.	.	.	.	.	.	.	26	8.7	3
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	22	52	36	52	78	201	105	192	221	437	47	9	.	.	.
	MEAN	4	10	7	13	20	50	26	48	55	87	9	2	.	.	.
BEARDED SEAL (H,S)	1981	.	.	.	.	.	.	.	.	.	21	0	0	21	7.0	3
	1982	0	0	2	0	1	0	0	6	0	3	0	0	12	1.0	12
	1983	0	0	0	0	3	6	11	1	0	11	2	0	34	2.8	12
	1984	0	0	2	14	6	2	3	2	9	11	0	0	49	4.1	12
	1985	0	0	0	0	5	4	0	3	1	2	0	0	15	1.3	12
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	0	0	4	14	15	12	14	12	10	48	2	0	.	.	.
	MEAN	0	0	1	4	4	3	4	3	3	10	0	0	.	.	.

For footnotes see Appendix 4.

Table 38. (cont'd)

HARBOUR SEAL (H,S)	1981	.	.	.	.	.	.	.	.	.	2	0	0	2	0.7	3
	1982	0	0	0	0	0	0	0	0	1	0	0	0	1	0.1	12
	1983	0	0	0	0	0	0	1	2	0	0	0	0	3	0.3	12
	1984	0	0	0	0	0	0	1	1	0	0	0	0	2	0.2	12
HARP SEAL (H,S)	1982	0	0	0	0	0	0	0	2	2	1	0	0	5	0.4	12
	1983	0	0	0	0	0	3	1	0	1	1	0	0	6	0.5	12
	1984	0	0	0	0	2	0	0	0	0	0	0	0	2	0.2	12
	1985	0	0	0	0	0	2	0	0	0	0	0	0	2	0.2	12
UNKNOWN SEAL	1983	0	0	0	0	0	0	1	0	0	0	0	0	1	0.1	12
WALRUS (H,S)	1985	0	0	0	0	1	0	0	0	0	0	0	0	1	0.1	12
BELUGA (H,A)	1982	0	0	0	0	0	0	9	74	2	0	0	0	85	7.1	12
	1983	0	0	0	0	0	0	8	48	2	0	0	0	58	4.8	12
	1984	0	0	0	0	0	0	35	15	0	0	0	0	50	4.2	12
	1985	0	0	0	0	0	0	10	84	0	0	0	0	94	7.8	12
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	0	0	0	0	0	0	62	221	4	0	0	0	.	.	.
	MEAN	0	0	0	0	0	0	16	55	1	0	0	0	.	.	.
CANADA GEESE (H,A)	1982	0	0	0	0	49	4	0	4	2	0	0	0	59	4.9	12
	1983	0	0	0	0	86	452	0	0	7	0	0	0	545	45.4	12
	1984	0	0	0	0	449	191	0	0	9	0	0	0	649	54.1	12
	1985	0	0	0	0	113	43	0	0	0	0	0	0	156	13.0	12
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	0	0	0	0	697	690	0	4	18	0	0	0	.	.	.
	MEAN	0	0	0	0	174	173	0	1	5	0	0	0	.	.	.
SNOW GEESE (H,A)	1982	0	0	0	0	667	38	0	0	9	0	0	0	714	59.5	12
	1983	0	0	0	0	0	90	0	0	3	0	0	0	93	7.8	12
	1984	0	0	0	0	108	14	0	0	1	0	0	0	123	10.3	12
	1985	0	0	0	0	518	187	0	0	0	0	0	0	705	58.8	12
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	0	0	0	0	1293	329	0	0	13	0	0	0	.	.	.
	MEAN	0	0	0	0	323	82	0	0	3	0	0	0	.	.	.
UNKNOWN GEESE	1984	0	0	0	0	12	0	0	0	0	0	0	0	12	1.0	12
	1985	0	0	0	0	24	6	0	0	0	0	0	0	30	2.5	12

For footnotes see Appendix 4.

Table 38. (cont'd)

OLD SQUAW	(L,S)	1983	0	0	0	0	0	8	0	0	0	0	0	0	8	0.7	12
		1984	0	0	0	0	0	8	0	0	0	0	0	0	8	0.7	12
EIDER	(L,S)	1981	.	.	.	.	.	.	.	.	.	2	0	0	2	0.7	3
		1983	0	0	0	0	0	3	0	0	0	0	0	0	3	0.3	12
		1984	0	0	0	0	1	2	0	0	9	6	0	1	19	1.6	12
		1985	0	0	0	0	0	2	0	0	0	2	0	0	4	0.3	12
		N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
		SUM	0	0	0	0	1	7	0	0	9	10	0	1	.	.	.
		MEAN	0	0	0	0	0	2	0	0	2	2	0	0	.	.	.
MALLARD	(L,S)	1982	0	0	0	0	0	2	0	0	0	0	0	0	2	0.2	12
		1984	0	0	0	0	0	0	0	0	0	0	1	0	1	0.1	12
PTARMIGAN	(L,A)	1981	.	.	.	.	.	.	.	.	.	75	13	3	91	30.3	3
		1982	14	18	15	48	26	0	12	12	32	0	1	13	191	15.9	12
		1983	4	7	0	2	39	16	6	11	16	9	68	12	190	15.8	12
		1984	5	0	20	111	7	0	0	1	134	45	0	0	323	26.9	12
		1985	2	0	0	40	72	8	0	0	0	0	0	0	122	10.2	12
		1986	0	76	24	.	.	.	.	.	.	.	.	.	100	33.3	3
		N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
		SUM	25	101	59	201	144	24	18	24	182	129	82	28	.	.	.
		MEAN	5	20	12	50	36	6	5	6	46	26	16	6	.	.	.
SNOWY OWL	(L,A)	1981	.	.	.	.	.	.	.	.	.	0	0	2	2	0.7	3
SWAN	(L,S)	1984	0	0	0	0	1	0	0	0	0	0	0	0	1	0.1	12
CANADA GOOSE EGGS		1984	0	0	0	0	384	0	0	0	0	0	0	0	384	32.0	12
SNOW GOOSE EGGS		1984	0	0	0	0	5	0	0	0	0	0	0	0	5	0.4	12
OLD SQUAW EGGS		1983	0	0	0	0	0	7	0	0	0	0	0	0	7	0.6	12

For footnotes see Appendix 4.

Table 38. (cont'd)

GOOSE EGGS	1983	0	0	0	0	0	1112	0	0	0	0	0	0	1112	92.7	12
	1984	0	0	0	0	0	61	0	0	0	0	0	0	61	5.1	12
	1985	0	0	0	0	141	204	0	0	0	0	0	0	345	28.8	12
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	0	0	0	0	141	1377	0	0	0	0	0	0	.	.	.
	MEAN	0	0	0	0	35	344	0	0	0	0	0	0	.	.	.
DUCK EGGS	1983	0	0	0	0	0	8	0	0	0	0	0	0	8	0.7	12
OTHER WATERFOWL EGGS	1983	0	0	0	0	0	7	1	0	0	0	0	0	8	0.7	12
ARCTIC CHARR (H,A)	1981	.	.	.	.	.	.	.	.	.	47	24	34	105	35.0	3
	1982	4	6	0	0	1	72	1080	1042	169	36	57	12	2479	206.6	12
	1983	3	0	0	0	61	364	562	694	258	85	38	30	2095	174.6	12
	1984	0	0	0	2	163	172	1249	608	152	78	109	68	2601	216.8	12
	1985	1	0	1	4	41	289	659	1779	357	59	119	93	3402	283.5	12
	1986	37	0	0	.	.	.	.	.	.	.	.	.	37	12.3	3
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	45	6	1	6	266	897	3550	4123	936	305	347	237	.	.	.
	MEAN	9	1	0	2	67	224	888	1031	234	61	69	47	.	.	.
LAKE TROUT (H,A)	1981	.	.	.	.	.	.	.	.	.	879	679	128	1686	562.0	3
	1982	91	12	79	182	237	38	9	65	75	28	5	0	821	68.4	12
	1983	48	11	213	115	281	72	0	24	130	68	159	231	1352	112.7	12
	1984	0	0	4	183	62	138	8	6	111	22	22	32	588	49.0	12
	1985	4	0	0	0	0	0	0	0	0	75	59	32	170	14.2	12
	1986	2	4	70	.	.	.	.	.	.	.	.	.	76	25.3	3
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	145	27	366	480	580	248	17	95	316	1072	924	423	.	.	.
	MEAN	29	5	73	120	145	62	4	24	79	214	185	85	.	.	.
WHITEFISH (H,S)	1981	.	.	.	.	.	.	.	.	.	0	27	356	383	127.7	3
	1982	13	0	20	248	424	57	0	0	23	0	0	0	785	65.4	12
	1983	0	0	0	0	0	0	0	0	0	0	101	8	109	9.1	12
	1984	0	0	0	0	0	0	0	0	45	0	0	2	47	3.9	12
	1985	0	0	0	0	0	0	0	0	0	0	68	52	120	10.0	12
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	13	0	20	248	424	57	0	0	68	0	196	418	.	.	.
	MEAN	3	0	4	62	106	14	0	0	17	0	39	84	.	.	.

For footnotes see Appendix 4.

Table 38. (cont'd)

ARCTIC GRAYLING (L,A)	1981	.	.	.	.	.	.	.	.	.	0	81	53	134	44.7	3
	1982	0	0	0	54	0	0	0	113	5	0	0	0	172	14.3	12
	1983	0	0	0	1	9	0	0	1	1	366	47	0	425	35.4	12
	1984	0	0	0	10	0	0	0	0	16	376	0	0	402	33.5	12
	1985	0	0	0	2	11	1	0	0	5	8	0	0	27	2.3	12
	1986	0	4	0	.	.	.	.	.	.	.	.	.	4	1.3	3
	N	5	5	5	4	4	4	4	4	4	5	5	5	.	.	.
	SUM	0	4	0	67	20	1	0	114	27	750	128	53	.	.	.
	MEAN	0	1	0	17	5	0	0	29	7	150	26	11	.	.	.
NORTHERN PIKE (L,S)	1981	.	.	.	.	.	.	.	.	.	0	2	0	2	0.7	3
	1982	9	0	0	0	0	0	0	0	0	0	0	0	9	0.8	12
	1983	0	0	0	0	0	0	0	52	35	0	0	0	87	7.3	12
	1984	0	0	0	0	0	0	0	0	16	0	0	0	16	1.3	12
	1985	0	0	0	0	0	1	1	0	0	0	0	0	2	0.2	12
FRESHWATER FISH	1983	0	0	0	0	0	0	0	0	2	0	19	0	21	1.8	12
	1984	0	0	0	0	0	0	0	0	1	0	7	2	10	0.8	12
	1985	0	0	0	0	1	1	0	0	51	0	0	0	53	4.4	12
ARCTIC COD (L,A)	1982	0	0	0	0	0	56	53	0	0	0	0	0	109	9.1	12
	1983	0	0	0	0	0	48	0	0	0	0	0	0	48	4.0	12
	1984	0	0	0	0	0	3	0	0	0	0	0	0	3	0.3	12
SCULPINS (L,A)	1983	0	0	0	0	0	1	0	0	0	0	0	0	1	0.1	12
	1984	0	0	0	0	0	3	0	0	0	0	0	0	3	0.3	12
SALTWATER FISH	1982	0	0	0	0	0	0	3	0	0	0	0	0	3	0.3	12
	1983	0	0	0	0	0	0	14	0	0	0	0	0	14	1.2	12

For footnotes see Appendix 4.

Table 39. The estimated harvest by Rankin Inlet hunters, expressed as number of animals, for the period November, 1981 to March, 1986. The actual harvest was used as the best estimate of the harvest for August, 1982 and June, 1983.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
CARIBOU KAMINURIAK (H,A)[6]	C[5]		1982	0[2]	0	0	0	0	0	0	2	53	7	2	10	74	6.2	12
			1983	0	0	1	0	0	0	0	0	0	0	2	1	4	0.3	12
			1984	0	4	0	0	0	0	0	2	0	0	0	0	6	0.5	12
			1985	0	0	0	1	0	0	0	4	1	3	5	0	14	1.2	12
			1986[1]	6	11	7	.	.	.	.	.	.	.	.	.	24	8.0	3
			N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
			SUM	6	15	8	1	0	0	0	8	54	10	9	11	.	.	.
			MEAN	1	3	2	0	0	0	0	2	14	3	2	2	.	.	.
		F	1981	.[3]	.	.	.	.	.	.	.	.	.	132	223	355	177.5	2
			1982	116	29	80	56	10	0	3	19	93	24	37	39	506	42.2	12
			1983	65	57	85	42	80	2	0	27	20	49	41	69	537	44.8	12
			1984	33	65	61	71	25	0	1	5	7	11	89	99	467	38.9	12
			1985	132	65	37	114	38	25	0	15	13	67	97	31	634	52.8	12
			1986	52	65	35	.	.	.	.	.	.	.	.	.	152	50.7	3
			N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
			SUM	398	281	298	283	153	27	4	66	133	151	396	461	.	.	.
			MEAN	80	56	60	71	38	7	1	17	33	38	79	92	.	.	.
		M	1981	.	.	.	.	.	.	.	.	.	.	43	94	137	68.5	2
			1982	28	114	107	163	127	7	74	163	320	91	37	29	1260	105.0	12
			1983	50	39	47	48	275	15	12	126	130	51	60	61	914	76.2	12
			1984	79	138	90	113	132	8	64	39	37	20	44	68	832	69.3	12
			1985	100	47	78	103	106	14	57	181	124	55	25	23	913	76.1	12
			1986	32	36	33	.	.	.	.	.	.	.	.	.	101	33.7	3
			N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
			SUM	289	374	355	427	640	44	207	509	611	217	209	275	.	.	.
			MEAN	58	75	71	107	160	11	52	127	153	54	42	55	.	.	.
		U	1982	0	21	3	0	0	0	0	0	0	36	11	1	72	6.0	12
			1983	0	0	2	0	0	0	9	2	25	6	0	26	70	5.8	12
			1984	0	2	22	14	19	13	1	1	0	0	14	0	86	7.2	12
			1985	9	0	0	0	0	12	4	4	0	0	0	4	33	2.8	12
			1986	3	1	1	.	.	.	.	.	.	.	.	.	5	1.7	3
			N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
			SUM	12	24	28	14	19	25	14	7	25	42	25	31	.	.	.
			MEAN	2	5	6	4	5	6	4	2	6	11	5	6	.	.	.

For footnotes see Appendix 4.

Table 39. (cont'd)

CARIBOU N. CHEST.	C	1986[4]	0	0	8	.	.	.	.	.	.	.	.	.	8	2.7	3
(H,A)	F	1984	0	21	0	0	0	0	0	0	19	0	0	0	40	3.3	12
		1986	0	0	25	.	.	.	.	.	.	.	.	.	25	8.3	3
	M	1984	0	15	0	0	0	0	0	1	29	0	0	0	45	3.8	12
		1985	0	60	0	3	0	0	4	0	0	0	0	0	67	5.6	12
		1986	0	0	29	.	.	.	.	.	.	.	.	.	29	9.7	3
	U	1984	0	0	0	0	0	0	0	7	0	0	0	0	7	0.6	12
		1986	0	0	2	.	.	.	.	.	.	.	.	.	2	0.7	3
MUSKOX (H,S)		1985	3	0	0	0	0	0	0	0	0	0	0	0	3	0.3	12
		1986	2	0	0	.	.	.	.	.	.	.	.	.	2	0.7	3
POLAR BEAR (H,S)		1981	.	.	.	.	.	.	.	.	.	.	3	2	5	2.5	2
		1982	0	0	1	3	0	0	0	0	0	0	5	0	9	0.8	12
		1983	0	1	1	0	12	0	0	0	0	0	3	3	20	1.7	12
		1984	0	1	1	0	0	1	0	0	0	0	1	0	4	0.3	12
		1985	0	2	0	0	6	3	0	0	0	0	10	0	21	1.8	12
		N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
		SUM	0	4	3	3	18	4	0	0	0	0	22	5	.	.	.
		MEAN	0	1	1	1	5	1	0	0	0	0	4	1	.	.	.
ARCTIC FOX (H,A)		1981	.	.	.	.	.	.	.	.	.	.	0	7	7	3.5	2
		1982	1	6	15	21	0	0	0	0	1	0	561	127	732	61.0	12
		1983	26	21	56	2	0	0	0	0	0	0	23	33	161	13.4	12
		1984	17	23	20	10	0	0	0	0	0	0	31	33	134	11.2	12
		1985	0	5	1	0	0	0	0	0	0	0	52	119	177	14.8	12
		1986	70	22	9	.	.	.	.	.	.	.	.	.	101	33.7	3
		N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
		SUM	114	77	101	33	0	0	0	0	1	0	667	319	.	.	.
		MEAN	23	15	20	8	0	0	0	0	0	0	133	64	.	.	.
WOLF (H,S)		1982	4	0	1	9	0	0	0	0	0	0	0	0	14	1.2	12
		1983	2	2	7	14	6	0	0	0	0	0	1	0	32	2.7	12
		1984	0	2	3	3	1	0	0	0	0	0	0	0	9	0.8	12
		1985	1	2	6	4	3	1	0	0	0	0	0	0	17	1.4	12
		1986	2	4	1	.	.	.	.	.	.	.	.	.	7	2.3	3
		N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
		SUM	9	10	18	30	10	1	0	0	0	0	1	0	.	.	.
		MEAN	2	2	4	8	3	0	0	0	0	0	0	0	.	.	.
WOLVERINE (H,S)		1983	0	0	0	0	9	0	0	0	0	0	0	0	9	0.8	12
		1984	0	1	0	0	0	0	0	0	0	0	0	0	1	0.1	12

For footnotes see Appendix 4.

Table 39. (cont'd)

ARCTIC HARE (L,S)	1981	.	.	.	.	.	.	.	.	.	.	.	.	.	.
	1982	0	0	0	0	0	0	0	0	0	0	0	2	2	1.0 2
	1983	5	0	2	0	0	0	0	0	7	0	0	0	7	0.6 12
	1984	0	0	0	0	4	0	0	0	0	0	0	0	7	0.6 12
	1985	0	0	0	0	0	0	0	0	0	0	1	0	5	0.4 12
	1986	2	2	0	.	.	.	.	.	8	2	0	0	10	0.8 12
														4	1.3 3
GROUND SQUIRREL (L,A)	1984	0	0	0	0	0	1	0	0	0	0	0	0	1	0.1 12
	1985	0	0	0	0	0	0	2	0	0	0	0	0	2	0.2 12
RINGED SEAL (H,A)	1981	.	.	.	.	.	.	.	.	.	.	.	.	.	.
	1982	0	9	17	33	22	101	159	77	34	71	0	0	0	0.0 2
	1983	11	16	9	10	18	74	79	68	51	49	34	8	565	47.1 12
	1984	4	0	6	4	48	125	116	28	11	77	23	0	408	34.0 12
	1985	11	2	10	10	22	73	114	30	22	66	16	19	454	37.8 12
	1986	3	9	14	.	.	.	.	.	.	.	25	2	387	32.3 12
	N	5	5	5	4	4	4	4	4	4	4	.	.	26	8.7 3
	SUM	29	36	56	57	110	373	468	203	118	263	5	5	.	. .
	MEAN	6	7	11	14	28	93	117	51	30	66	98	29	.	. .
												20	6	.	. .
BEARDED SEAL (H,S)	1982	0	0	0	3	0	2	2	5	1	6	2	0	21	1.8 12
	1983	0	0	0	4	0	1	3	0	2	2	1	0	13	1.1 12
	1984	0	0	2	4	3	2	3	1	0	11	0	0	26	2.2 12
	1985	0	0	0	0	0	3	8	6	5	8	0	0	30	2.5 12
	1986	0	0	1	.	.	.	.	.	.	.	.	.	1	0.3 3
	N	5	5	5	4	4	4	4	4	4	4	5	5	.	. .
	SUM	0	0	3	11	3	8	16	12	8	27	3	0	.	. .
	MEAN	0	0	1	3	1	2	4	3	2	7	1	0	.	. .
HARBOUR SEAL (H,S)	1983	0	0	0	0	0	0	0	0	2	0	0	0	2	0.2 12
	1984	0	0	0	0	0	0	0	0	1	2	0	0	3	0.3 12
	1985	0	0	0	0	0	1	0	0	0	2	0	0	3	0.3 12
HARP SEAL (H,S)	1984	0	0	0	0	0	0	0	1	0	0	0	0	1	0.1 12
	1985	0	0	0	0	0	0	0	0	1	1	0	0	2	0.2 12
UNKNOWN SEAL	1984	0	0	0	0	4	0	0	0	0	0	3	0	7	0.6 12

For footnotes see Appendix 4.

Table 39. (cont'd)

WALRUS (H,S)	1982	0	0	0	0	0	2	0	0	0	0	0	0	2	0.2	12
	1983	0	0	2	0	36	0	0	9	0	0	0	0	47	3.9	12
	1984	0	0	0	1	0	0	0	0	0	0	0	0	1	0.1	12
	1985	0	2	0	0	1	0	2	0	0	0	0	0	5	0.4	12
	1986	0	0	2	.	.	.	.	.	.	.	.	.	2	0.7	3
BELUGA (H,A)	1982	0	0	0	0	0	0	6	17	11	0	0	0	34	2.8	12
	1983	0	0	0	0	0	0	0	29	0	0	0	0	29	2.4	12
	1984	0	0	0	0	0	2	13	49	5	0	0	0	69	5.8	12
	1985	0	0	0	0	0	0	4	37	6	0	0	0	47	3.9	12
	N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
	SUM	0	0	0	0	0	2	23	132	22	0	0	0	.	.	.
	MEAN	0	0	0	0	0	1	6	33	6	0	0	0	.	.	.
CANADA GEESE (L,S)	1982	0	0	0	0	76	1112	0	0	0	0	0	0	1188	99.0	12
	1983	0	0	0	0	6	14	0	0	0	0	0	0	20	1.7	12
	1984	0	0	0	0	16	376	0	9	0	0	0	0	401	33.4	12
	1985	0	0	0	0	65	247	2	1	0	0	0	0	315	26.3	12
	N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
	SUM	0	0	0	0	163	1749	2	10	0	0	0	0	.	.	.
	MEAN	0	0	0	0	41	437	1	3	0	0	0	0	.	.	.
SNOW GEESE (L,S)	1982	0	0	0	9	0	0	2	0	41	0	0	0	52	4.3	12
	1983	0	0	0	0	0	80	0	0	18	0	0	0	98	8.2	12
	1984	0	0	0	0	209	51	0	0	41	0	0	0	301	25.1	12
	1985	0	0	0	0	184	335	0	0	0	0	0	0	519	43.3	12
	N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
	SUM	0	0	0	9	393	466	2	0	100	0	0	0	.	.	.
	MEAN	0	0	0	2	98	117	1	0	25	0	0	0	.	.	.
BRANT GEESE (L,S)	1984	0	0	0	0	0	0	11	0	0	0	0	0	11	0.9	12
UNKNOWN GEESE	1982	0	0	0	0	0	0	0	0	1	0	0	0	1	0.1	12
	1985	0	0	0	0	0	3	0	0	0	0	0	0	3	0.3	12
SANDHILL CRANE (L,S)	1982	0	0	0	0	10	0	0	0	0	0	0	0	10	0.8	12
	1984	0	0	0	0	3	0	0	0	0	0	0	0	3	0.3	12

For footnotes see Appendix 4.

Table 39. (cont'd)

EIDER (L,S)	1982	0	6	3	0	0	0	0	0	23	0	0	0	32	2.7	12
	1983	0	0	0	0	6	0	0	0	0	0	1	0	7	0.6	12
	1984	0	0	1	0	0	2	14	7	3	6	0	4	37	3.1	12
	1985	0	5	0	0	0	0	0	0	0	8	0	0	13	1.1	12
	N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
	SUM	0	11	4	0	6	2	14	7	26	14	1	4	.	.	.
	MEAN	0	2	1	0	2	1	4	2	7	4	0	1	.	.	.
PTARMIGAN (L,S)	1982	0	0	0	15	22	0	0	0	11	13	3	0	64	5.3	12
	1983	2	0	0	0	201	0	0	9	0	0	0	0	212	17.7	12
	1984	0	12	0	155	97	13	0	7	7	22	7	0	320	26.7	12
	1985	0	5	0	11	82	55	0	0	2	4	0	0	159	13.3	12
	N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
	SUM	2	17	0	181	402	68	0	16	20	39	10	0	.	.	.
	MEAN	0	3	0	45	101	17	0	4	5	10	2	0	.	.	.
UNKNOWN FOWL	1984	0	0	0	0	0	0	1	0	0	0	0	0	1	0.1	12
	1985	0	0	0	0	0	0	0	6	1	0	0	0	7	0.6	12
SWAN (L,S)	1984	0	0	0	0	6	1	0	2	0	0	0	0	9	0.8	12
CANADA GOOSE EGGS	1984	0	0	0	0	0	94	0	0	0	0	0	0	94	7.8	12
	1985	0	0	0	0	0	13	0	0	0	0	0	0	13	1.1	12
GOOSE EGGS	1985	0	0	0	0	0	83	137	0	0	0	0	0	220	18.3	12
OTHER WATERFOWL EGGS	1984	0	0	0	0	0	0	22	0	0	0	0	0	22	1.8	12
ARCTIC CHARR (H,A)	1981	.	.	.	.	.	.	.	.	.	.	0	719	719	359.5	2
	1982	0	353	0	134	82	913	7366	1386	115	791	727	359	12226	1018.8	12
	1983	214	0	0	0	0	420	788	2117	65	102	443	492	4641	386.8	12
	1984	333	86	93	57	74	861	724	1804	45	70	763	205	5115	426.3	12
	1985	124	44	96	114	139	1290	1448	2873	0	404	615	316	7463	621.9	12
	1986	315	114	107	.	.	.	.	.	.	.	.	.	536	178.7	3
	N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
	SUM	986	597	296	305	295	3484	10326	8180	225	1367	2548	2091	.	.	.
	MEAN	197	119	59	76	74	871	2582	2045	56	342	510	418	.	.	.

For footnotes see Appendix 4.

Table 39. (cont'd)

LANDLOCKED ARCTIC CHARR (L,A)	1985	0	0	0	0	0	0	0	0	0	32	0	0	32	2.7	12
LAKE TROUT (H,A)	1981	.	.	.	.	.	.	.	.	.	.	0	33	33	16.5	2
	1982	0	15	78	59	0	0	0	0	0	0	45	0	197	16.4	12
	1983	0	20	0	0	289	0	0	0	0	0	54	0	363	30.3	12
	1984	0	20	0	120	232	0	10	21	0	0	103	0	506	42.2	12
	1985	1	16	14	0	173	127	0	0	15	0	0	2	348	29.0	12
	1986	16	0	0	.	.	.	.	.	.	.	.	.	16	5.3	3
	N	5	5	5	4	4	4	4	4	4	4	5	5	.	.	.
	SUM	17	71	92	179	694	127	10	21	15	0	202	35	.	.	.
	MEAN	3	14	18	45	174	32	3	5	4	0	40	7	.	.	.
WHITEFISH (H,S)	1983	0	0	0	0	0	0	0	0	0	0	7	0	7	0.6	12
	1984	0	0	0	1	0	0	0	0	0	0	8	0	9	0.8	12
ARCTIC GRAYLING (L,A)	1982	0	0	0	0	0	0	0	0	10	0	0	0	10	0.8	12
	1984	0	0	0	0	0	0	0	0	0	0	49	0	49	4.1	12
FRESHWATER FISH	1982	0	0	0	30	117	0	0	0	0	0	0	0	147	12.3	12
	1983	0	0	0	104	0	0	0	0	0	0	0	0	104	8.7	12
ARCTIC COD (L,A)	1985	0	0	0	0	0	0	12	0	0	0	0	0	12	1.0	12
SCULPINS (L,A)	1985	0	0	0	0	0	0	13	0	0	0	0	0	13	1.1	12
SALTWATER FISH	1983	0	0	0	52	0	0	0	0	0	0	0	0	52	4.3	12

For footnotes see Appendix 4.

Table 40. The estimated harvest by Repulse Bay hunters, expressed as number of animals for the period October, 1981 to March, 1986. Data were not available for December, 1982. The best estimate of the harvest for December 1981, March 1982, and April 1983 was the actual harvest.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	$\bar{x}$	NUM
CARIBOU	KAMINURIAK	F[5]	1984[4]	0[2]	0	0	0	6	0	0	0	0	0	0	0	6	0.5	12
	(H,S)[6]	M	1982	3	0	0	0	0	0	0	0	0	0	0	.	3	0.3	11
			1983	0	0	0	0	0	2	0	0	0	0	0	0	2	0.2	12
			1984	0	0	0	0	34	0	0	0	0	0	0	0	34	2.8	12
		U	1984	0	0	0	0	1	0	0	0	0	0	0	0	1	0.1	12
CARIBOU	WAGER	C	1982	10	0	0	0	0	0	9	24	9	0	0	.	52	4.7	11
	(H,A)		1983	0	0	0	0	0	0	0	7	6	0	3	0	16	1.3	12
			1984	0	0	0	0	0	0	0	4	0	0	0	1	5	0.4	12
			1985	0	0	0	0	0	0	0	21	30	3	4	1	59	4.9	12
			1986[1]	2	3	10	.	.	.	.	.	.	.	.	.	15	5.0	3
			N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
			SUM	12	3	10	0	0	0	9	56	45	3	7	2	.	.	.
			MEAN	2	1	2	0	0	0	2	14	11	1	1	1	.	.	.
		F	1981	.[3]	.	.	.	.	.	.	.	.	66	49	22	137	45.7	3
			1982	46	74	17	44	10	0	3	90	25	24	53	.	386	35.1	11
			1983	41	32	36	28	9	8	3	29	15	24	11	17	253	21.1	12
			1984	20	47	60	20	6	9	7	79	36	25	72	38	419	34.9	12
			1985	48	79	72	55	13	2	4	24	61	25	22	12	417	34.8	12
			1986	59	42	56	.	.	.	.	.	.	.	.	.	157	52.3	3
			N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
			SUM	214	274	241	147	38	19	17	222	137	164	207	89	.	.	.
			MEAN	43	55	48	37	10	5	4	56	34	33	41	22	.	.	.
		M	1981	.	.	.	.	.	.	.	.	.	97	51	19	167	55.7	3
			1982	34	30	25	49	42	141	22	192	88	16	46	.	685	62.3	11
			1983	70	27	43	33	41	62	63	68	81	52	9	17	566	47.2	12
			1984	15	65	39	34	30	91	59	194	94	69	49	29	768	64.0	12
			1985	18	46	48	48	75	44	125	121	141	51	44	31	792	66.0	12
			1986	44	83	31	.	.	.	.	.	.	.	.	.	158	52.7	3
			N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
			SUM	181	251	186	164	188	338	269	575	404	285	199	96	.	.	.
			MEAN	36	50	37	41	47	85	67	144	101	57	40	24	.	.	.
		U	1982	0	5	14	3	0	8	22	6	0	0	0	.	58	5.3	11
			1983	0	0	6	0	0	0	0	0	0	19	14	0	39	3.3	12
			1984	5	12	37	46	0	10	5	1	16	4	1	3	140	11.7	12
			1985	1	0	4	0	0	15	7	15	4	0	0	0	46	3.8	12
			1986	0	0	4	.	.	.	.	.	.	.	.	.	4	1.3	3
			N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
			SUM	6	17	65	49	0	33	34	22	20	23	15	3	.	.	.
			MEAN	1	3	13	12	0	8	9	6	5	5	3	1	.	.	.

For footnotes see Appendix 4.

Table 40. (cont'd)

CARIBOU SOUTHAMPTON (H,S)	U	1985	0	0	0	0	0	0	0	0	1	0	0	0	1	0.1 12
CARIBOU BEVERLY	F	1981	.	.	.	.	.	.	.	.	.	5	2	0	7	2.3 3
(H,S)	M	1981	.	.	.	.	.	.	.	.	.	3	0	2	5	1.7 3
CARIBOU N. CHEST. (H,S)	M	1984	0	0	0	0	23	0	0	0	0	0	0	0	23	1.9 12
CARIBOU OTHER	M	1984	0	0	0	0	7	0	0	0	0	0	0	0	7	0.6 12
	U	1984	0	0	0	0	1	0	0	0	0	0	0	0	1	0.1 12
MUSKOX (H,S)		1985	0	1	0	0	0	0	0	0	0	0	0	0	1	0.1 12
POLAR BEAR (H,S)		1981	.	.	.	.	.	.	.	.	.	0	15	1	16	5.3 3
		1982	0	0	0	0	0	0	0	0	0	0	15	.	15	1.4 11
		1983	0	1	1	1	0	0	0	0	0	0	8	2	13	1.1 12
		1984	0	2	2	0	0	0	0	0	0	0	4	1	9	0.8 12
		1985	0	3	0	0	0	0	0	0	0	0	8	0	11	0.9 12
		1986	2	4	0	.	.	.	.	.	.	.	.	.	6	2.0 3
		N	5	5	5	4	4	4	4	4	4	5	5	4	.	. .
		SUM	2	10	3	1	0	0	0	0	0	0	50	4	.	. .
		MEAN	0	2	1	0	0	0	0	0	0	0	10	1	.	. .
BLACK BEAR (L,S)		1983	0	1	0	0	0	0	0	0	0	0	0	0	1	0.1 12
GRIZZLY BEAR (H,S)		1982	0	0	0	0	0	5	0	0	0	0	0	.	5	0.5 11
		N	5	5	5	4	4	4	4	4	4	5	5	4	.	. .
		SUM	0	0	0	0	0	5	0	0	0	0	0	0	.	. .
		MEAN	0	0	0	0	0	1	0	0	0	0	0	0	.	. .

For footnotes see Appendix 4.

Table 40. (cont'd)

ARCTIC FOX (H,A)	1981	.	.	.	.	.	.	.	.	.	0	3	0	3	1.0	3
	1982	13	19	8	34	0	0	0	0	0	0	7	.	81	7.4	11
	1983	37	22	22	16	0	0	0	0	0	0	97	78	272	22.7	12
	1984	35	20	34	16	0	0	0	0	0	1	337	77	520	43.3	12
	1985	30	35	24	4	0	0	0	0	0	0	2	0	95	7.9	12
	1986	0	1	5	.	.	.	.	.	.	.	.	.	6	2.0	3
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	115	97	93	70	0	0	0	0	0	1	446	155	.	.	.
	MEAN	23	19	19	18	0	0	0	0	0	0	89	39	.	.	.
RED FOX (H,S)	1982	0	0	1	0	0	0	0	0	0	0	0	.	1	0.1	11
	1983	0	0	0	0	0	0	0	0	0	0	1	0	1	0.1	12
	1984	0	2	0	0	0	0	0	0	0	0	21	0	23	1.9	12
	1985	1	0	3	0	0	0	0	0	0	0	4	6	14	1.2	12
	1986	0	1	0	.	.	.	.	.	.	.	.	.	1	0.3	3
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	1	3	4	0	0	0	0	0	0	0	26	6	.	.	.
	MEAN	0	1	1	0	0	0	0	0	0	0	5	2	.	.	.
WOLF (H,S)	1982	5	18	4	2	13	0	0	0	0	0	0	.	42	3.8	11
	1983	6	0	6	4	3	0	0	0	0	4	10	2	35	2.9	12
	1984	2	4	18	2	4	0	0	0	0	1	8	0	39	3.3	12
	1985	2	10	9	10	4	3	0	0	0	0	6	8	52	4.3	12
	1986	4	13	5	.	.	.	.	.	.	.	.	.	22	7.3	3
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	19	45	42	18	24	3	0	0	0	5	24	10	.	.	.
	MEAN	4	9	8	5	6	1	0	0	0	1	5	3	.	.	.
WOLVERINE (H,S)	1982	0	0	3	0	0	0	0	0	0	0	0	.	3	0.3	11
	1983	0	0	0	0	0	0	0	0	0	0	3	0	3	0.3	12
	1984	0	0	4	2	1	0	0	0	0	0	0	1	8	0.7	12
	1985	0	0	0	0	0	0	0	0	0	0	0	1	1	0.1	12
ARCTIC HARE (L,S)	1981	.	.	.	.	.	.	.	.	.	13	0	0	13	4.3	3
	1982	0	5	0	0	0	0	0	0	2	0	0	.	7	0.6	11
	1983	0	0	1	1	5	0	0	0	0	0	0	0	7	0.6	12
	1984	0	4	0	2	0	0	0	0	0	0	0	1	7	0.6	12
	1985	0	0	1	1	0	0	0	0	7	0	0	0	9	0.8	12
	1986	0	0	2	.	.	.	.	.	.	.	.	.	2	0.7	3
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	0	9	4	4	5	0	0	0	9	13	0	1	.	.	.
	MEAN	0	2	1	1	1	0	0	0	2	3	0	0	.	.	.

For footnotes see Appendix 4.

Table 40. (cont'd)

RINGED SEAL (H,A)	1981	.	.	.	.	.	.	.	.	.	357	10	4	371	123.7	3
	1982	0	0	2	29	6	172	71	114	45	20	3	.	462	42.0	11
	1983	0	8	14	10	12	134	30	69	45	87	14	0	423	35.3	12
	1984	8	10	9	16	19	122	126	85	57	123	17	0	592	49.3	12
	1985	14	0	4	18	180	90	145	64	195	62	22	6	800	66.7	12
	1986	3	21	12	.	.	.	.	.	.	.	.	.	36	12.0	3
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	25	39	41	73	217	518	372	332	342	649	66	10	.	.	.
	MEAN	5	8	8	18	54	130	93	83	86	130	13	3	.	.	.
BEARDED SEAL (H,S)	1981	.	.	.	.	.	.	.	.	.	2	0	5	7	2.3	3
	1982	0	0	0	0	0	0	0	12	2	0	0	.	14	1.3	11
	1983	0	0	0	0	0	0	1	9	6	2	0	0	18	1.5	12
	1984	0	0	0	0	0	0	5	16	2	1	0	0	24	2.0	12
	1985	0	0	0	0	0	0	0	10	4	3	0	1	18	1.5	12
	1986	0	0	1	.	.	.	.	.	.	.	.	.	1	0.3	3
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	0	0	1	0	0	0	6	47	14	8	0	6	.	.	.
	MEAN	0	0	0	0	0	0	2	12	4	2	0	2	.	.	.
HARP SEAL (H,S)	1982	0	0	0	0	0	0	0	3	0	0	0	.	3	0.3	11
	1984	0	0	0	0	0	0	2	4	0	1	0	0	7	0.6	12
	1985	0	0	10	1	0	0	15	3	0	0	0	0	29	2.4	12
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	0	0	10	1	0	0	17	10	0	1	0	0	.	.	.
	MEAN	0	0	2	0	0	0	4	3	0	0	0	0	.	.	.
WALRUS (H,A)	1981	.	.	.	.	.	.	.	.	.	16	0	0	16	5.3	3
	1982	0	0	0	0	0	0	0	0	5	0	0	.	5	0.5	11
	1983	0	0	0	0	0	5	4	0	0	4	0	0	13	1.1	12
	1984	0	0	0	0	0	0	0	3	2	0	0	0	5	0.4	12
	1985	0	0	0	0	0	0	1	0	17	4	0	0	22	1.8	12
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	0	0	0	0	0	5	5	3	24	24	0	0	.	.	.
	MEAN	0	0	0	0	0	1	1	1	6	5	0	0	.	.	.
BELUGA (H,A)	1981	.	.	.	.	.	.	.	.	.	3	0	0	3	1.0	3
	1982	0	0	0	0	0	0	0	15	20	0	0	.	35	3.2	11
	1983	0	0	0	0	0	0	8	16	17	0	0	0	41	3.4	12
	1984	0	0	0	0	0	0	2	12	11	0	0	0	25	2.1	12
	1985	0	0	0	0	0	0	1	3	0	0	0	0	4	0.3	12
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	0	0	0	0	0	0	11	46	48	3	0	0	.	.	.
	MEAN	0	0	0	0	0	0	3	12	12	1	0	0	.	.	.
NARWHAL (H,A)	1982	0	0	0	0	0	0	3	6	0	0	0	.	9	0.8	11
	1983	0	0	0	0	0	0	4	2	0	0	0	0	6	0.5	12
	1984	0	0	0	0	0	0	14	12	2	0	0	0	28	2.3	12
	1985	0	0	0	0	0	0	14	7	0	0	5	0	26	2.2	12
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	0	0	0	0	0	0	35	27	2	0	5	0	.	.	.
	MEAN	0	0	0	0	0	0	9	7	1	0	1	0	.	.	.

For footnotes see Appendix 4.

Table 40. (cont'd)

CANADA GEESE (L,S)	1983	0	0	0	0	0	0	0	0	2	0	0	0	2	0.2 12
	1984	0	0	0	0	4	3	0	0	0	0	0	0	7	0.6 12
	1985	0	0	0	0	7	12	1	0	3	0	0	0	23	1.9 12
SNOW GEESE (L,S)	1982	0	0	0	0	19	8	0	0	0	0	0	.	27	2.5 11
	1984	0	0	0	0	4	0	0	0	0	0	0	0	4	0.3 12
ROSS'S GEESE (L,S)	1982	0	0	0	0	3	5	0	0	0	0	0	.	8	0.7 11
	1983	0	0	0	0	0	0	10	0	0	0	0	0	10	0.8 12
UNKNOWN GEESE	1985	0	0	0	0	4	0	0	0	0	0	0	0	4	0.3 12
SANDHILL CRANE (L,S)	1984	0	0	0	0	1	0	0	0	0	0	0	0	1	0.1 12
OLD SQUAW (L,S)	1984	0	0	0	0	0	6	0	0	0	0	0	0	6	0.5 12
GUILLEMOT (L,S)	1982	0	0	0	0	0	0	6	3	0	0	0	.	9	0.8 11
EIDER (L,S)	1982	0	0	0	0	0	0	9	3	0	0	0	.	12	1.1 11
	1983	0	0	0	0	0	15	7	0	0	0	0	0	22	1.8 12
	1984	0	0	0	0	0	5	0	0	0	0	0	0	5	0.4 12
	1985	0	0	0	0	0	0	4	0	0	0	0	0	4	0.3 12

For footnotes see Appendix 4.

Table 40. (cont'd)

PTARMIGAN (L,S)	1982	0	0	0	0	96	29	0	0	117	10	0	.	252	22.9	11
	1983	0	0	0	0	3	0	0	0	0	0	0	0	3	0.3	12
	1984	0	6	0	5	0	4	0	6	61	0	4	0	86	7.2	12
	1985	2	0	0	0	20	5	1	0	26	13	0	0	67	5.6	12
	1986	3	0	0	.	.	.	.	.	.	.	.	.	3	1.0	3
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	5	6	0	5	119	38	1	6	204	23	4	0	.	.	.
	MEAN	1	1	0	1	30	10	0	2	51	5	1	0	.	.	.
UNKNOWN FOWL	1983	0	0	0	0	0	0	7	0	0	0	0	0	7	0.6	12
ARCTIC CHARR (H,A)	1981	.	.	.	.	.	.	.	.	.	133	289	283	705	235.0	3
	1982	0	0	23	0	157	313	47	390	29	542	418	.	1919	174.5	11
	1983	0	0	14	0	0	63	76	109	2	147	564	52	1027	85.6	12
	1984	0	0	0	0	14	655	265	493	9	631	426	350	2843	236.9	12
	1985	260	0	0	30	195	501	1017	749	0	27	639	0	3418	284.8	12
	1986	0	0	63	.	.	.	.	.	.	.	.	.	63	21.0	3
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	260	0	100	30	366	1532	1405	1741	40	1480	2336	685	.	.	.
	MEAN	52	0	20	8	92	383	351	435	10	296	467	171	.	.	.
LAKE TROUT (H,A)	1981	.	.	.	.	.	.	.	.	.	745	5	0	750	250.0	3
	1982	75	0	1	15	199	345	0	0	11	0	5	.	651	59.2	11
	1983	0	0	8	46	8	2	0	0	0	2	0	0	66	5.5	12
	1984	0	0	0	0	60	0	0	0	0	0	225	0	285	23.8	12
	1985	0	0	0	0	68	115	8	0	0	16	0	1	208	17.3	12
	1986	0	0	4	.	.	.	.	.	.	.	.	.	4	1.3	3
	N	5	5	5	4	4	4	4	4	4	5	5	4	.	.	.
	SUM	75	0	13	61	335	462	8	0	11	763	235	1	.	.	.
	MEAN	15	0	3	15	84	116	2	0	3	153	47	0	.	.	.
ARCTIC GRAYLING (L,A)	1982	0	0	0	0	0	0	0	0	14	0	0	.	14	1.3	11
FRESHWATER FISH	1983	0	0	0	0	0	0	0	0	0	216	0	0	216	18.0	12

For footnotes see Appendix 4.

Table 41. The estimated harvest by Whale Cove hunters, expressed as number of animals for the period October 1981 to March 1983[3] and November 1983 to March 1986. The best estimate of the harvest for October, 1981, 1982, 1983, and 1984, November 1982 and 1983, and December 1983 was the actual reported harvest.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
CARIBOU KAMINURIAK	C[5]		1982	0[2]	0	0	1	0	0	0	13	36	0	2	0	52	4.3	12
(H,A)[6]			1984	0	0	0	0	0	0	0	0	0	0	4	1	5	0.4	12
			1985	7	1	0	0	0	1	1	1	1	0	0	2	14	1.2	12
			1986	9	4	4	.	.	.	.	.	.	.	.	.	17	5.7	3
			N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
			SUM	16	5	4	1	0	1	1	14	37	0	6	3	.	.	.
			MEAN	3	1	1	0	0	0	0	5	12	0	1	1	.	.	.
		F	1981	.[3]	.	.	.	.	.	.	.	.	30	71	91	192	64.0	3
			1982	39	88	58	57	27	0	4	5	34	0	29	54	395	32.9	12
			1983	18	32	65	.	.	.	.	.	.	.	36	15	166	33.2	5
			1984	10	32	77	38	11	10	0	0	13	6	33	58	288	24.0	12
			1985	97	44	91	46	20	3	5	13	0	32	29	33	413	34.4	12
			1986	73	43	41	.	.	.	.	.	.	.	.	.	157	52.3	3
			N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
			SUM	237	239	332	141	58	13	9	18	47	68	198	251	.	.	.
			MEAN	47	48	66	47	19	4	3	6	16	17	40	50	.	.	.
		M	1981	.	.	.	.	.	.	.	.	.	12	16	12	40	13.3	3
			1982	29	42	44	52	65	0	19	97	101	17	23	34	523	43.6	12
			1983	47	11	15	.	.	.	.	.	.	.	28	10	111	22.2	5
			1984	52	46	52	16	22	18	12	17	21	22	17	6	301	25.1	12
			1985	11	11	20	7	52	28	46	45	42	30	14	10	316	26.3	12
			1986	20	6	7	.	.	.	.	.	.	.	.	.	33	11.0	3
			N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
			SUM	159	116	138	75	139	46	77	159	164	81	98	72	.	.	.
			MEAN	32	23	28	25	46	15	26	53	55	20	20	14	.	.	.
		U	1982	0	0	4	0	0	0	0	25	0	0	0	6	35	2.9	12
			1983	5	0	20	.	.	.	.	.	.	.	7	2	34	6.8	5
			1984	0	0	0	0	0	0	0	0	0	0	0	1	1	0.1	12
			1985	0	0	0	9	1	0	2	0	0	0	0	0	12	1.0	12
			1986	3	0	1	.	.	.	.	.	.	.	.	.	4	1.3	3
			N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
			SUM	8	0	25	9	1	0	2	25	0	0	7	9	.	.	.
			MEAN	2	0	5	3	0	0	1	8	0	0	1	2	.	.	.

For footnotes see Appendix 4.

Table 47. (cont'd)

CARIBOU WAGER (H,S)	M	1981[4]	.	.	.	.	.	.	.	.	.	4	0	0	4	1.3	3
MUSKOX (H,S)		1985	0	2	0	0	0	0	0	0	0	0	0	0	2	0.2	12
POLAR BEAR (H,A)		1981	.	.	.	.	.	.	.	.	.	0	4	2	6	2.0	3
		1982	0	0	0	1	0	0	0	0	0	1	2	2	6	0.5	12
		1983	0	0	0	.	.	.	.	.	.	.	5	0	5	1.0	5
		1984	0	0	2	0	1	0	0	0	0	0	7	0	10	0.8	12
		1985	0	0	0	0	1	0	0	0	0	0	10	0	11	0.9	12
		N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
		SUM	0	0	2	1	2	0	0	0	0	1	28	4	.	.	.
		MEAN	0	0	0	0	1	0	0	0	0	0	6	1	.	.	.
BLACK BEAR (L,S)		1982	0	0	0	0	0	0	0	0	1	0	0	0	1	0.1	12
ARCTIC FOX (H,A)		1981	.	.	.	.	.	.	.	.	.	0	0	0	0	0.0	3
		1982	0	1	0	4	0	0	0	0	0	0	94	80	179	14.9	12
		1983	32	8	30	.	.	.	.	.	.	.	10	26	106	21.2	5
		1984	0	0	0	0	0	0	0	0	0	0	120	4	124	10.3	12
		1985	0	1	0	0	0	0	0	0	0	0	0	2	3	0.3	12
		1986	2	0	0	.	.	.	.	.	.	.	.	.	2	0.7	3
		N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
		SUM	34	10	30	4	0	0	0	0	0	0	224	112	.	.	.
		MEAN	7	2	6	1	0	0	0	0	0	0	45	22	.	.	.
RED FOX (H,S)		1983	2	0	0	.	.	.	.	.	.	.	0	0	2	0.4	5
		1985	0	0	0	0	0	0	0	0	0	0	0	1	1	0.1	12
WOLF (H,S)		1982	0	0	2	3	0	0	0	0	0	0	0	0	5	0.4	12
		1983	0	0	0	.	.	.	.	.	.	.	1	3	4	0.8	5
		1984	0	3	0	2	0	0	0	0	0	1	0	0	6	0.5	12
		1985	0	2	12	0	0	0	0	0	0	0	0	0	14	1.2	12
ARCTIC HARE (L,S)		1981	.	.	.	.	.	.	.	.	.	9	1	0	10	3.3	3
		1982	4	0	0	0	0	0	0	0	0	0	0	0	4	0.3	12
		1983	2	0	5	.	.	.	.	.	.	.	0	0	7	1.4	5
		1984	0	0	0	0	0	0	0	3	5	1	0	0	9	0.8	12
		1985	1	1	3	1	0	0	4	0	2	0	1	1	14	1.2	12
		1986	0	2	0	.	.	.	.	.	.	.	.	.	2	0.7	3
		N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
		SUM	7	3	8	1	0	0	4	3	7	10	2	1	.	.	.
		MEAN	1	1	2	0	0	0	1	1	2	3	0	0	.	.	.

For footnotes see Appendix 4.

Table 41. (cont'd)

RINGED SEAL (H,A)	1981	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
	1982	0	5	17	8	29	20	3	31	8	8	2	11	142	11.8	12
	1983	10	11	8	.	.	.	.	.	.	.	3	0	32	6.4	5
	1984	6	5	4	5	21	29	13	6	14	29	40	28	200	16.7	12
	1985	9	11	16	9	34	17	36	16	1	10	1	6	166	13.8	12
	1986	7	3	4	.	.	.	.	.	.	.	.	.	14	4.7	3
	N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
	SUM	32	35	49	22	84	66	52	53	23	47	50	45	.	.	.
	MEAN	6	7	10	7	28	22	17	18	8	12	10	9	.	.	.
BEARDED SEAL (H,S)	1982	0	1	0	4	0	0	2	0	0	2	0	0	9	0.8	12
	1984	0	0	0	1	0	7	0	2	0	17	0	0	27	2.3	12
	1985	0	1	1	0	0	0	2	0	3	0	0	0	7	0.6	12
HARBOUR SEAL (H,S)	1982	0	0	2	0	0	0	0	0	0	0	0	0	2	0.2	12
	1983	0	0	3	.	.	.	.	.	.	.	0	0	3	0.6	5
	1984	0	0	0	0	0	0	3	3	0	0	0	0	6	0.5	12
HARP SEAL (H,S)	1982	0	0	0	0	0	0	0	0	1	0	0	0	1	0.1	12
	1983	0	0	3	.	.	.	.	.	.	.	0	0	3	0.6	5
WALRUS (H,S)	1981	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
	1982	0	1	0	1	4	0	0	0	0	1	0	0	1	0.3	3
											0	0	0	6	0.5	12
BELUGA (H,A)	1982	0	0	0	0	0	0	0	1	2	0	0	0	3	0.3	12
	1984	0	0	0	0	0	0	0	19	5	0	0	0	24	2.0	12
	1985	0	0	0	0	0	0	0	17	2	0	0	0	19	1.6	12
	N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
	SUM	0	0	0	0	0	0	0	37	9	0	0	0	.	.	.
	MEAN	0	0	0	0	0	0	0	12	3	0	0	0	.	.	.
NARWHAL (H,S)	1982	0	0	0	0	0	0	0	1	0	0	0	0	1	0.1	12
CANADA GEESE (L,S)	1982	0	0	0	0	71	29	0	0	0	0	0	0	100	8.3	12
	1984	0	0	0	0	11	13	0	0	0	0	0	0	24	2.0	12
	1985	0	0	0	0	68	12	19	0	5	0	0	0	104	8.7	12
	N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
	SUM	0	0	0	0	150	54	19	0	5	0	0	0	.	.	.
	MEAN	0	0	0	0	50	18	6	0	2	0	0	0	.	.	.

For footnotes see Appendix 4.

Table 41. (cont'd)

SNOW GEESE (L,S)	1982	0	0	0	0	109	40	0	0	0	0	0	0	149	12.4	12
	1984	0	0	0	0	226	273	37	0	4	0	0	0	540	45.0	12
	1985	0	0	0	0	55	0	0	0	29	0	0	0	84	7.0	12
	N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
	SUM	0	0	0	0	390	313	37	0	33	0	0	0	.	.	.
	MEAN	0	0	0	0	130	104	12	0	11	0	0	0	.	.	.
ROSS'S GEESE (L,S)	1982	0	0	0	0	0	2	0	0	0	0	0	0	2	0.2	12
UNKNOWN GEESE	1985	0	0	0	0	338	197	1	0	62	0	0	0	598	49.8	12
EIDER (L,S)	1981	.	.	.	.	.	.	.	.	.	1	0	0	1	0.3	3
	1982	0	0	0	0	8	0	0	0	0	0	0	0	8	0.7	12
	1984	0	0	0	0	9	0	0	0	0	8	0	0	17	1.4	12
	1985	0	3	0	0	0	1	0	0	0	0	0	0	4	0.3	12
PTARMIGAN (L,A)	1981	.	.	.	.	.	.	.	.	.	9	0	0	9	3.0	3
	1982	0	0	0	0	6	0	0	0	2	13	0	0	21	1.8	12
	1983	0	9	0	.	.	.	.	.	.	.	0	0	9	1.8	5
	1984	0	0	.	5	7	0	0	0	0	0	0	0	12	1.1	11
	1985	0	10	0	0	0	0	0	0	0	0	0	0	10	0.8	12
CANADA GOOSE EGGS	1985	0	0	0	0	0	0	2	0	0	0	0	0	2	0.2	12
GOOSE EGGS	1984	0	0	0	0	24	0	0	0	0	0	0	0	24	2.0	12
	1985	0	0	0	0	124	599	0	0	0	0	0	0	723	60.3	12
ARCTIC CHARR (H,A)	1981	.	.	.	.	.	.	.	.	.	42	11	166	219	73.0	3
	1982	111	42	79	58	86	332	630	6583	44	36	69	13	8083	673.6	12
	1983	2	25	0	.	.	.	.	.	.	.	63	30	120	24.0	5
	1984	23	0	0	7	3	88	261	467	20	195	117	16	1197	99.8	12
	1985	4	5	0	0	56	76	117	416	3	10	9	6	702	58.5	12
	1986	40	20	3	.	.	.	.	.	.	.	.	.	63	21.0	3
	N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
	SUM	180	92	82	65	145	496	1008	7466	67	283	269	231	.	.	.
	MEAN	36	18	16	22	48	165	336	2489	22	71	54	46	.	.	.

For footnotes see Appendix 4.

Table 41. (cont'd)

LAKE TROUT (H,A)	1981	.	.	.	.	.	.	.	.	.	4	33	156	193	64.3	3
	1982	59	12	14	71	52	15	23	120	3	15	26	58	468	39.0	12
	1983	11	56	18	.	.	.	.	.	.	.	0	0	85	17.0	5
	1984	17	14	92	93	80	18	0	0	0	0	9	0	323	26.9	12
	1985	0	0	0	6	52	44	7	1	0	3	0	0	113	9.4	12
	1986	10	9	0	.	.	.	.	.	.	.	.	.	19	6.3	3
	N	5	5	5	3	3	3	3	3	3	4	5	5	.	.	.
	SUM	97	91	124	170	184	77	30	121	3	22	68	214	.	.	.
	MEAN	19	18	25	57	61	26	10	40	1	6	14	43	.	.	.
WHITEFISH (H,S)	1982	0	0	0	0	0	0	0	0	0	18	15	43	76	6.3	12
ARCTIC GRAYLING (L,A)	1981	.	.	.	.	.	.	.	.	.	0	0	2	2	0.7	3
	1985	0	0	0	0	5	0	0	0	0	0	0	0	5	0.4	12
NORTHERN PIKE (L,S)	1982	0	0	0	0	2	0	0	0	0	0	0	0	2	0.2	12
FRESHWATER FISH	1982	0	0	3	8	0	0	0	0	0	0	0	0	11	0.9	12
SALTWATER FISH	1982	0	0	0	0	6	0	0	0	0	0	0	0	6	0.5	12

For footnotes see Appendix 4.

Table 42. The estimated harvest of Kaminuriak caribou by hunters from the seven Keewatin communities expressed as numbers of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
BAKER LAKE	1981	. [1]	.	.	.	.	.	.	.	.	.	295	266	561	280.5 [3]	2
	1982	608	462	330	241	278	10	453	.	496	.	239	110	3227	322.7	10
	1983	286	0	71	334	161	0	241	216	412	92	41	79	1933	193.3	10
	1984	0	0	0	93	0	51	311	63	33	0	2	0	553	92.2	6
	1985	0	0	226	112	87	0	78	0	165	171	155	78	1072	134.0	8
	1986	0	0	81	.	.	.	.	.	.	.	.	.	81	81.0	1
CORAL HARBOUR	1984	0	0	4	0	0	0	0	0	0	0	0	0	4	4.0	1
	1985	0	0	0	9	0	0	0	0	0	0	0	0	9	9.0	1
CHESTERFIELD INLET	1981	.	.	.	.	.	.	.	.	.	.	.	.	0	0.0	0
	1982	4	17	.	.	.	.	.	6	1	0	27	0	55	11.0	5
	1983	17	16	0	0	0	3	0	5	2	0	1	17	61	8.7	7
	1984	31	0	1	0	40	0	0	8	0	0	0	26	106	21.2	5
	1985	0	9	30	3	2	0	2	0	0	11	10	0	67	9.6	7
ESKIMO POINT	1981	.	.	.	.	.	.	.	.	.	455	539	156	1150	383.3	3
	1982	185	216	258	296	70	10	314	530	730	289	121	103	3122	260.2	12
	1983	189	135	310	236	146	30	144	238	313	365	112	153	2371	197.6	12
	1984	193	355	465	336	33	20	173	113	460	128	134	155	2565	213.8	12
	1985	166	211	143	228	123	83	150	152	273	500	124	108	2261	188.4	12
	1986	124	141	400	.	.	.	.	.	.	.	.	.	665	221.7	3
REPULSE BAY	1982	3	0 [2]	0	0	0	0	0	0	0	0	0	.	3	3.0	1
	1983	0	0	0	0	0	2	0	0	0	0	0	0	2	2.0	1
	1984	0	0	0	0	41	0	0	0	0	0	0	0	41	41.0	1
RANKIN INLET	1981	.	.	.	.	.	.	.	.	.	.	175	317	492	246.0	2
	1982	144	164	190	219	137	7	77	184	486	158	87	79	1912	159.3	12
	1983	115	96	135	90	355	17	21	155	175	106	103	157	1525	127.1	12
	1984	112	209	173	198	176	21	66	47	44	31	147	167	1391	115.9	12
	1985	241	112	115	218	144	51	61	204	138	125	127	58	1594	132.8	12
	1986	93	113	76	.	.	.	.	.	.	.	.	.	282	94.0	3
WHALE COVE	1981	.	.	.	.	.	.	.	.	.	42	87	103	232	77.3	3
	1982	68	130	106	110	92	0	23	140	171	17	54	94	1005	91.4	11
	1983	70	43	100	.	.	.	.	.	.	.	71	27	311	62.2	5
	1984	62	78	129	54	33	28	12	17	34	28	54	66	595	49.6	12
	1985	115	56	111	62	73	32	54	59	43	62	43	45	755	62.9	12
	1986	105	53	53	.	.	.	.	.	.	.	.	.	211	70.3	3
KEEWATIN	1981	.	.	.	.	.	.	.	.	.	497	1096	842	2435	811.7	3
	1982	1012	989	884	866	577	27	867	860	1864	464	528	386	9324	777.0	12
	1983	677	290	616	660	662	52	406	614	902	563	328	433	6203	516.9	12
	1984	398	642	772	681	323	120	562	248	571	187	337	414	5255	437.9	12
	1985	522	388	625	632	429	166	345	415	619	869	459	289	5758	479.8	12
	1986	322	307	610	.	.	.	.	.	.	.	.	.	1239	413.0	3

For footnotes see Appendix 4.

Table 43. The estimated harvest of Beverly caribou by Baker Lake and Repulse Bay hunters expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
BAKER LAKE	1981	[1] .	.	.	.	.	.	.	.	.	.	.	[2] .	.	[3] .	.
	1982	11	32	44	56	9	0	10	.	19	.	102	0	102	102.0	1
	1983	0	362	251	258	319	0	184	47	92	151	41	0	222	27.8	8
	1984	251	593	749	658	60	0	77	0	437	569	467	175	2306	230.6	10
	1985	460	382	67	30	0	0	0	0	363	341	0	340	3734	414.9	9
REPULSE BAY	1981	.	.	.	.	.	.	.	.	.	8	126	0	1769	252.7	7
KEEWATIN	1981	.	.	.	.	.	.	.	.	.	8	2	2	12	4.0	3
	1982	11	32	44	56	9	0	10	.	19	.	104	2	114	38.0	3
	1983	0	362	251	258	319	0	184	47	92	151	41	0	222	27.8	8
	1984	251	593	749	658	60	0	77	0	437	569	467	175	2306	230.6	10
	1985	460	382	67	30	0	0	0	0	363	341	126	0	1769	252.7	7

For footnotes see Appendix 4.

Table 44. The estimated harvest of Wager Bay caribou by Baker Lake, Coral Harbour, Eskimo Point, Repulse Bay and Whale Cove hunters, expressed as numbers of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
BAKER LAKE	1982	0	0	0	0	0	0	0	.[1]	6	.	0[2]	0	6	6.0[3]	1
	1983	0	0	0	0	0	401	192	248	480	31	8	68	1428	204.0	7
	1984	2	0	0	4	508	154	134	511	505	631	279	35	2763	276.3	10
	1985	0	15	0	97	90	134	237	560	848	207	35	287	2510	251.0	10
	1986	349	424	418	.	.	.	.	.	.	.	.	.	1191	397.0	3
CORAL HARBOUR	1984	0	0	0	193	0	0	0	12	0	0	0	0	205	102.5	2
	1985	0	0	0	84	126	0	0	0	0	0	0	0	210	105.0	2
ESKIMO POINT	1983	0	0	0	0	0	0	0	0	1	0	0	0	1	1.0	1
REPULSE BAY	1981	.	.	.	.	.	.	.	.	.	163	100	41	304	101.3	3
	1982	90	109	56	96	52	149	56	312	122	40	99	.	1181	107.4	11
	1983	111	59	85	61	50	70	66	104	102	95	37	34	874	72.8	12
	1984	40	124	136	100	36	110	71	278	146	98	122	71	1332	111.0	12
	1985	67	125	124	103	88	61	136	181	236	79	70	44	1314	109.5	12
	1986	105	128	101	.	.	.	.	.	.	.	.	.	334	111.3	3
WHALE COVE	1981	.	.	.	.	.	.	.	.	.	4	0	0	4	4.0	1
KEEWATIN	1981	.	.	.	.	.	.	.	.	.	167	100	41	308	102.7	3
	1982	90	109	56	96	52	149	56	312	128	40	99	0	1187	107.9	11
	1983	111	59	85	61	50	471	258	352	583	126	45	102	2303	191.9	12
	1984	42	124	136	297	544	264	205	801	651	729	401	106	4300	358.3	12
	1985	67	140	124	284	304	195	373	741	1084	286	105	331	4034	336.2	12
	1986	454	552	519	.	.	.	.	.	.	.	.	.	1525	508.3	3

For footnotes see Appendix 4.

Table 45. The estimated harvest of polar bears by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{x}$	NUM
BAKER LAKE	1985	0	0[2]	1	0	0	0	0	0	0	0	0	0	1	1.0[3]	1
CORAL HARBOUR	1982	.[1]	0	.	.	.	2	0	0	0	1	6	5	14	3.5	4
	1983	0	0	0	0	.	.	.	4	2	20	9	1	36	7.2	5
	1984	4	0	0	0	0	0	0	0	0	32	9	0	45	15.0	3
	1985	1	0	3	4	4	0	0	0	0	11	35	3	61	8.7	7
	1986	2	1	2	.	.	.	.	.	.	.	.	.	5	1.7	3
CHESTERFIELD INLET	1982	3	0	.	.	.	.	.	0	0	0	0	1	4	2.0	2
	1983	2	0	1	7	0	0	0	0	0	0	4	0	14	3.5	4
	1984	1	0	4	0	0	0	0	0	0	0	0	0	5	2.5	2
	1985	0	1	2	0	2	0	0	0	0	1	2	0	8	1.6	5
ESKIMO POINT	1981	.	.	.	.	.	.	.	.	.	0	10	0	10	10.0	1
	1982	0	0	0	4	0	0	0	0	0	0	13	0	17	8.5	2
	1983	0	0	1	1	0	0	0	0	0	7	12	0	21	5.3	4
	1984	0	0	1	0	1	0	0	0	0	0	13	0	15	5.0	3
	1985	0	0	0	0	0	0	0	0	0	0	8	0	8	8.0	1
	1986	0	0	3	.	.	.	.	.	.	.	.	.	3	3.0	1
REPULSE BAY	1981	.	.	.	.	.	.	.	.	.	0	15	1	16	8.0	2
	1982	0	0	0	0	0	0	0	0	0	0	15	.	15	15.0	1
	1983	0	1	1	1	0	0	0	0	0	0	8	2	13	2.6	5
	1984	0	2	2	0	0	0	0	0	0	0	4	1	9	2.3	4
	1985	0	3	0	0	0	0	0	0	0	0	8	0	11	5.5	2
	1986	2	4	0	.	.	.	.	.	.	.	.	.	6	3.0	2
RANKIN INLET	1981	.	.	.	.	.	.	.	.	.	.	3	2	5	2.5	2
	1982	0	0	1	3	0	0	0	0	0	0	5	0	9	3.0	3
	1983	0	1	1	0	12	0	0	0	0	0	3	3	20	4.0	5
	1984	0	1	1	0	0	1	0	0	0	0	1	0	4	1.0	4
	1985	0	2	0	0	6	3	0	0	0	0	10	0	21	5.3	4
WHALE COVE	1981	.	.	.	.	.	.	.	.	.	0	4	2	6	3.0	2
	1982	0	0	0	1	0	0	0	0	0	1	2	2	6	1.5	4
	1983	0	0	0	.	.	.	.	.	.	.	5	0	5	5.0	1
	1984	0	0	2	0	1	0	0	0	0	0	7	0	10	3.3	3
	1985	0	0	0	0	1	0	0	0	0	0	10	0	11	5.5	2
KEEWATIN	1981	.	.	.	.	.	.	.	.	.	0	32	5	37	18.5	2
	1982	3	0	1	8	0	2	0	0	0	2	41	8	65	9.3	7
	1983	2	2	4	9	12	0	0	4	2	27	41	6	109	10.9	10
	1984	5	3	10	0	2	1	0	0	0	32	34	1	88	11.0	8
	1985	1	6	6	4	13	3	0	0	0	12	73	3	121	13.4	9
	1986	4	5	5	.	.	.	.	.	.	.	.	.	14	4.7	3

For footnotes see Appendix 4.

Table 46. The estimated harvest of Arctic fox by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{x}$	NUM
BAKER LAKE	1981	.[1]	.	.	.	.	.	.	.	.	.	13	0[2]	13	13.0[3]	1
	1982	0	50	71	39	0	0	0	.	0	.	417	149	726	145.2	5
	1983	30	7	0	0	0	0	0	0	0	0	17	275	329	82.3	4
	1984	124	156	180	5	0	0	0	0	0	0	526	1090	2081	346.8	6
	1985	758	585	216	113	0	0	0	0	0	0	497	250	2419	403.2	6
	1986	65	152	75	.	.	.	.	.	.	.	.	.	292	97.3	3
CORAL HARBOUR	1982	.	50	.	.	.	0	0	0	0	166	167	103	486	121.5	4
	1983	117	160	224	17	.	.	.	23	21	0	126	51	739	92.4	8
	1984	85	55	89	123	0	0	0	0	0	0	109	123	584	97.3	6
	1985	29	115	76	74	12	0	0	0	0	0	17	69	392	56.0	7
	1986	13	2	35	.	.	.	.	.	.	.	.	.	50	16.7	3
CHESTERFIELD INLET	1982	10	15	.	.	.	.	.	0	0	0	223	92	340	85.0	4
	1983	85	73	34	69	0	0	0	0	0	0	0	26	287	57.4	5
	1984	2	3	2	2	0	0	0	0	0	24	8	18	59	8.4	7
	1985	0	5	6	0	0	0	0	0	0	0	26	0	37	12.3	3
	1986	0	0	4	.	.	.	.	.	.	.	.	.	4	4.0	1
ESKIMO POINT	1981	.	.	.	.	.	.	.	.	.	0	137	90	227	113.5	2
	1982	40	58	48	172	0	0	0	0	0	24	1487	469	2298	328.3	7
	1983	210	62	80	27	6	0	0	0	0	0	40	103	528	75.4	7
	1984	117	142	163	55	1	0	0	0	0	0	227	263	968	138.3	7
	1985	28	51	262	186	0	0	0	0	0	8	255	176	966	138.0	7
	1986	40	120	40	.	.	.	.	.	.	.	.	.	200	66.7	3
REPULSE BAY	1981	.	.	.	.	.	.	.	.	.	0	3	0	3	3.0	1
	1982	13	19	8	34	0	0	0	0	0	0	7	.	81	16.2	5
	1983	37	22	22	16	0	0	0	0	0	0	97	78	272	45.3	6
	1984	35	20	34	16	0	0	0	0	0	1	337	77	520	74.3	7
	1985	30	35	24	4	0	0	0	0	0	0	2	0	95	19.0	5
	1986	0	1	5	.	.	.	.	.	.	.	.	.	6	3.0	2
RANKIN INLET	1981	.	.	.	.	.	.	.	.	.	.	0	7	7	7.0	1
	1982	1	6	15	21	0	0	0	0	1	0	561	127	732	104.6	7
	1983	26	21	56	2	0	0	0	0	0	0	23	33	161	26.8	6
	1984	17	23	20	10	0	0	0	0	0	0	31	33	134	22.3	6
	1985	0	5	1	0	0	0	0	0	0	0	52	119	177	44.3	4
	1986	70	22	9	.	.	.	.	.	.	.	.	.	101	33.7	3
WHALE COVE	1981	.	.	.	.	.	.	.	.	.	0	0	0	0	0.0	0
	1982	0	1	0	4	0	0	0	0	0	0	94	80	179	44.8	4
	1983	32	8	30	.	.	.	.	.	.	.	10	26	106	21.2	5
	1984	0	0	0	0	0	0	0	0	0	0	120	4	124	62.0	2
	1985	0	1	0	0	0	0	0	0	0	0	0	2	3	1.5	2
	1986	2	0	0	.	.	.	.	.	.	.	.	.	2	2.0	1
KEEWATIN	1981	.	.	.	.	.	.	.	.	.	0	153	97	250	125.0	2
	1982	64	199	142	270	0	0	0	0	1	190	2956	1020	4842	605.3	8
	1983	537	353	446	131	6	0	0	23	21	0	313	592	2422	269.1	9
	1984	380	399	488	211	1	0	0	0	0	25	1358	1608	4470	558.8	8
	1985	845	797	585	377	12	0	0	0	0	8	849	616	4089	511.1	8
	1986	190	297	168	.	.	.	.	.	.	.	.	.	655	218.3	3

For footnotes see Appendix 4.

Table 47. The estimated harvest of wolves by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
BAKER LAKE	1982	4	0	16	2	0	0	0	.[1]	0	.	6	0[2]	28	7.0[3]	4
	1983	0	0	3	0	2	0	0	0	0	1	3	11	20	4.0	5
	1984	0	12	25	1	0	0	0	0	0	0	11	9	58	11.6	5
	1985	7	16	18	1	0	0	0	0	0	0	0	21	63	12.6	5
	1986	3	5	13	.	.	.	.	.	.	.	.	.	21	7.0	3
CORAL HARBOUR	1984	0	0	0	0	1	0	0	0	0	0	0	0	1	1.0	1
	1985	0	0	0	4	4	0	0	0	0	2	0	0	10	3.3	3
CHESTERFIELD INLET	1982	0	0	.	.	.	.	.	0	0	0	0	4	4	4.0	1
	1983	0	2	0	3	1	0	0	0	0	0	2	9	17	3.4	5
	1984	6	5	0	0	0	0	0	0	0	0	0	4	15	5.0	3
	1985	0	4	0	0	0	0	0	0	0	0	2	7	13	4.3	3
	1986	0	3	2	.	.	.	.	.	.	.	.	.	5	2.5	2
ESKIMO POINT	0	0	0	0	0	0	0	0	0	0	0	0	10	10	10.0	1
	1982	0	0	2	22	0	0	0	0	0	0	1	0	25	8.3	3
	1983	0	0	0	12	8	0	0	0	0	2	4	5	31	6.2	5
	1984	0	2	12	30	2	0	0	0	0	0	3	1	50	8.3	6
	1985	0	0	3	32	2	0	0	0	0	0	4	10	51	10.2	5
	1986	6	2	53	.	.	.	.	.	.	.	.	.	61	20.3	3
REPULSE BAY	1982	5	18	4	2	13	0	0	0	0	0	0	.	42	8.4	5
	1983	6	0	6	4	3	0	0	0	0	4	10	2	35	5.0	7
	1984	2	4	18	2	4	0	0	0	0	1	8	0	39	5.6	7
	1985	2	10	9	10	4	3	0	0	0	0	6	8	52	6.5	8
	1986	4	13	5	.	.	.	.	.	.	.	.	.	22	7.3	3
RANKIN INLET	1982	4	0	1	9	0	0	0	0	0	0	0	0	14	4.7	3
	1983	2	2	7	14	6	0	0	0	0	0	1	0	32	5.3	6
	1984	0	2	3	3	1	0	0	0	0	0	0	0	9	2.3	4
	1985	1	2	6	4	3	1	0	0	0	0	0	0	17	2.8	6
	1986	2	4	1	.	.	.	.	.	.	.	.	.	7	2.3	3
WHALE COVE	1982	0	0	2	3	0	0	0	0	0	0	0	0	5	2.5	2
	1983	0	0	0	.	.	.	.	.	.	.	1	3	4	2.0	2
	1984	0	3	0	2	0	0	0	0	0	1	0	0	6	2.0	3
	1985	0	2	12	0	0	0	0	0	0	0	0	0	14	7.0	2
KEEWATIN	0	0	0	0	0	0	0	0	0	0	0	0	10	10	10.0	1
	1982	13	18	25	38	13	0	0	0	0	0	7	4	118	16.9	7
	1983	8	4	16	33	20	0	0	0	0	7	21	30	139	17.4	8
	1984	8	28	58	38	8	0	0	0	0	2	22	14	178	22.3	8
	1985	10	34	48	51	13	4	0	0	0	2	12	46	220	24.4	9
	1986	15	27	74	.	.	.	.	.	.	.	.	.	116	38.7	3

For footnotes see Appendix 4.

Table 48. The estimated harvest of ringed seals by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
BAKER LAKE	1983	0	0[2]	0	0	1	0	0	0	0	0	0	0	1	1.0[3]	1
	1984	0	0	0	0	2	3	0	0	1	0	0	0	6	2.0	3
CORAL HARBOUR	1982	.	10	.[1]	.	.	278	107	108	52	42	30	52	679	84.9	8
	1983	67	62	40	22	.	.	.	192	49	59	87	13	591	65.7	9
	1984	134	101	55	41	57	142	74	20	45	46	20	21	756	63.0	12
	1985	89	93	19	22	30	113	83	18	33	79	86	74	739	61.6	12
	1986	81	83	78	.	.	.	.	.	.	.	.	.	242	80.7	3
CHESTERFIELD INLET	1982	7	6	.	.	.	.	.	4	29	44	2	1	93	13.3	7
	1983	2	2	7	19	8	16	20	6	11	6	5	2	104	8.7	12
	1984	0	0	3	0	7	6	4	3	7	7	0	0	37	5.3	7
	1985	1	0	7	0	6	9	2	6	2	14	10	0	57	6.3	9
	1986	9	6	14	.	.	.	.	.	.	.	.	.	29	9.7	3
ESKIMO POINT	1981	.	.	.	.	.	.	.	.	.	108	2	3	113	37.7	3
	1982	1	4	3	10	5	37	20	149	70	70	4	6	379	31.6	12
	1983	3	4	0	2	2	82	31	27	12	110	36	0	309	30.9	10
	1984	17	29	20	39	36	40	44	13	132	133	0	0	503	50.3	10
	1985	0	3	0	1	35	42	10	3	7	16	5	0	122	13.6	9
	1986	1	12	13	.	.	.	.	.	.	.	.	.	26	8.7	3
REPULSE BAY	1981	.	.	.	.	.	.	.	.	.	357	10	4	371	123.7	3
	1982	0	0	2	29	6	172	71	114	45	20	3	.	462	51.3	9
	1983	0	8	14	10	12	134	30	69	45	87	14	0	423	42.3	10
	1984	8	10	9	16	19	122	126	85	57	123	17	0	592	53.8	11
	1985	14	0	4	18	180	90	145	64	195	62	22	6	800	72.7	11
	1986	3	21	12	.	.	.	.	.	.	.	.	.	36	12.0	3
RANKIN INLET	1981	.	.	.	.	.	.	.	.	.	.	0	0	0	0.0	0
	1982	0	9	17	33	22	101	159	77	34	71	34	8	565	51.4	11
	1983	11	16	9	10	18	74	79	68	51	49	23	0	408	37.1	11
	1984	4	0	6	4	48	125	116	28	11	77	16	19	454	41.3	11
	1985	11	2	10	10	22	73	114	30	22	66	25	2	387	32.3	12
	1986	3	9	14	.	.	.	.	.	.	.	.	.	26	8.7	3
WHALE COVE	1981	.	.	.	.	.	.	.	.	.	0	4	0	4	4.0	1
	1982	0	5	17	8	29	20	3	31	8	8	2	11	142	12.9	11
	1983	10	11	8	.	.	.	.	.	.	.	3	0	32	8.0	4
	1984	6	5	4	5	21	29	13	6	14	29	40	28	200	16.7	12
	1985	9	11	16	9	34	17	36	16	1	10	1	6	166	13.8	12
	1986	7	3	4	.	.	.	.	.	.	.	.	.	14	4.7	3
KEEWATIN	1981	.	.	.	.	.	.	.	.	.	465	16	7	488	162.7	3
	1982	8	34	39	80	62	608	360	483	238	255	75	78	2320	193.3	12
	1983	93	103	78	63	41	306	160	362	168	311	168	15	1868	155.7	12
	1984	169	145	97	105	190	467	377	155	267	415	93	68	2548	212.3	12
	1985	124	109	56	60	307	344	390	137	260	247	149	88	2271	189.3	12
	1986	104	134	135	.	.	.	.	.	.	.	.	.	373	124.3	3

For footnotes see Appendix 4.

Table 49. The estimated harvest of bearded seals by hunters from Chesterfield Inlet, Coral Harbour, Eskimo Point, Rankin Inlet, Repulse Bay and Whale Cove expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
CORAL HARBOUR	1982	.[1]	16	.	.	.	2	3	3	10	1	0[2]	0	35	5.8[3]	6
	1983	0	0	3	6	.	.	.	20	33	2	4	1	69	9.9	7
	1984	4	7	17	2	0	5	12	4	10	2	0	3	66	6.6	10
	1985	6	5	8	2	0	0	6	14	6	2	0	0	49	6.1	8
	1986	5	6	0	.	.	.	.	.	.	.	.	.	11	5.5	2
CHESTERFIELD INLET	1982	0	0	.	.	.	.	.	0	2	0	0	0	2	2.0	1
	1983	0	0	0	0	0	0	0	0	0	3	0	0	3	3.0	1
	1984	0	0	0	0	0	0	0	0	1	0	0	0	1	1.0	1
	1985	1	0	0	1	1	1	0	0	0	1	0	0	5	1.0	5
	1986	0	0	3	.	.	.	.	.	.	.	.	.	3	3.0	1
ESKIMO POINT	1981	.	.	.	.	.	.	.	.	.	21	0	0	21	21.0	1
	1982	0	0	2	0	1	0	0	6	0	3	0	0	12	3.0	4
	1983	0	0	0	0	3	6	11	1	0	11	2	0	34	5.7	6
	1984	0	0	2	14	6	2	3	2	9	11	0	0	49	6.1	8
	1985	0	0	0	0	5	4	0	3	1	2	0	0	15	3.0	5
REPULSE BAY	1981	.	.	.	.	.	.	.	.	.	2	0	5	7	3.5	2
	1982	0	0	0	0	0	0	0	12	2	0	0	.	14	7.0	2
	1983	0	0	0	0	0	0	1	9	6	2	0	0	18	4.5	4
	1984	0	0	0	0	0	0	5	16	2	1	0	0	24	6.0	4
	1985	0	0	0	0	0	0	0	10	4	3	0	1	18	4.5	4
	1986	0	0	1	.	.	.	.	.	.	.	.	.	1	1.0	1
RANKIN INLET	1982	0	0	0	3	0	2	2	5	1	6	2	0	21	3.0	7
	1983	0	0	0	4	0	1	3	0	2	2	1	0	13	2.2	6
	1984	0	0	2	4	3	2	3	1	0	11	0	0	26	3.7	7
	1985	0	0	0	0	0	3	8	6	5	8	0	0	30	6.0	5
	1986	0	0	1	.	.	.	.	.	.	.	.	.	1	1.0	1
WHALE COVE	1982	0	1	0	4	0	0	2	0	0	2	0	0	9	2.3	4
	1984	0	0	0	1	0	7	0	2	0	17	0	0	27	6.8	4
	1985	0	1	1	0	0	0	2	0	3	0	0	0	7	1.8	4
KEEWATIN	1981	.	.	.	.	.	.	.	.	.	23	0	5	28	14.0	2
	1982	0	17	2	7	1	4	7	26	15	12	2	0	93	9.3	10
	1983	0	0	3	10	3	7	15	30	41	20	7	1	137	13.7	10
	1984	4	7	21	21	9	16	23	25	22	42	0	3	193	17.5	11
	1985	7	6	9	3	6	8	16	33	19	16	0	1	124	11.3	11
	1986	5	6	5	.	.	.	.	.	.	.	.	.	16	5.3	3

For footnotes see Appendix 4.

Table 50. The estimated harvest of beluga whales by hunters from Chesterfield Inlet, Coral Harbour, Eskimo Point, Rankin Inlet, Repulse Bay and Whale Cove expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{X}$ [3]	NUM
CORAL HARBOUR	1982	.[1]	0[2]	.	.	.	0	17	75	32	0	0	1	125	31.3	4
	1983	0	0	0	0	.	.	.	77	48	2	1	0	128	32.0	4
	1984	2	1	0	0	0	3	33	50	24	6	0	0	121	17.3	7
	1985	0	0	0	0	0	0	15	60	13	0	0	0	88	29.3	3
CHESTERFIELD INLET	1981	.	.	.	.	.	.	.	.	.	0	0	0	0	0.0	0
	1982	0	0	.	.	.	.	.	5	3	0	0	0	8	4.0	2
	1983	0	0	0	0	0	0	0	7	0	0	0	0	7	7.0	1
	1984	0	0	0	0	0	0	4	4	4	0	0	0	12	4.0	3
	1985	0	0	0	0	0	0	0	16	16	0	0	0	32	16.0	2
ESKIMO POINT	1982	0	0	0	0	0	0	9	74	2	0	0	0	85	28.3	3
	1983	0	0	0	0	0	0	8	48	2	0	0	0	58	19.3	3
	1984	0	0	0	0	0	0	35	15	0	0	0	0	50	25.0	2
	1985	0	0	0	0	0	0	10	84	0	0	0	0	94	47.0	2
REPULSE BAY	1981	.	.	.	.	.	.	.	.	.	3	0	0	3	3.0	1
	1982	0	0	0	0	0	0	0	15	20	0	0	.	35	17.5	2
	1983	0	0	0	0	0	0	8	16	17	0	0	0	41	13.7	3
	1984	0	0	0	0	0	0	2	12	11	0	0	0	25	8.3	3
	1985	0	0	0	0	0	0	1	3	0	0	0	0	4	2.0	2
RANKIN INLET	1982	0	0	0	0	0	0	6	17	11	0	0	0	34	11.3	3
	1983	0	0	0	0	0	0	0	29	0	0	0	0	29	29.0	1
	1984	0	0	0	0	0	2	13	49	5	0	0	0	69	17.3	4
	1985	0	0	0	0	0	0	4	37	6	0	0	0	47	15.7	3
WHALE COVE	1982	0	0	0	0	0	0	0	1	2	0	0	0	3	1.5	2
	1984	0	0	0	0	0	0	0	19	5	0	0	0	24	12.0	2
	1985	0	0	0	0	0	0	0	17	2	0	0	0	19	9.5	2
KEEWATIN	1981	.	.	.	.	.	.	.	.	.	3	0	0	3	3.0	1
	1982	0	0	0	0	0	0	32	187	70	0	0	1	290	72.5	4
	1983	0	0	0	0	0	0	16	177	67	2	1	0	263	52.6	5
	1984	2	1	0	0	0	5	87	149	49	8	0	0	301	43.0	7
	1985	0	0	0	0	0	0	30	217	37	0	0	0	284	94.7	3

For footnotes see Appendix 4.

Table 51. The estimated harvest of Canada Geese by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{x}$	NUM
BAKER LAKE	1984	0	0	0[2]	0	154	142	0	0	0	0	0	0	296	148.0[3]	2
	1985	0	0	0	0	479	144	0	0	0	0	0	0	623	311.5	2
CORAL HARBOUR	1982	.[1]	0	.	.	.	41	407	199	0	0	0	9	656	164.0	4
	1983	0	0	0	0	.	.	.	9	10	0	0	0	19	9.5	2
	1984	0	0	0	0	13	101	0	23	0	0	0	0	137	45.7	3
	1985	0	0	0	0	30	25	0	0	0	0	0	0	55	27.5	2
CHESTERFIELD INLET	1984	0	0	0	0	0	8	0	0	0	0	0	0	8	8.0	1
	1985	0	0	0	0	0	30	0	0	0	0	0	0	30	30.0	1
ESKIMO POINT	1982	0	0	0	0	49	4	0	4	2	0	0	0	59	14.8	4
	1983	0	0	0	0	86	452	0	0	7	0	0	0	545	181.7	3
	1984	0	0	0	0	449	191	0	0	9	0	0	0	649	216.3	3
	1985	0	0	0	0	113	43	0	0	0	0	0	0	156	78.0	2
REPULSE BAY	1983	0	0	0	0	0	0	0	0	2	0	0	0	2	2.0	1
	1984	0	0	0	0	4	3	0	0	0	0	0	0	7	3.5	2
	1985	0	0	0	0	7	12	1	0	3	0	0	0	23	5.8	4
RANKIN INLET	1982	0	0	0	0	76	1112	0	0	0	0	0	0	1188	594.0	2
	1983	0	0	0	0	6	14	0	0	0	0	0	0	20	10.0	2
	1984	0	0	0	0	16	376	0	9	0	0	0	0	401	133.7	3
	1985	0	0	0	0	65	247	2	1	0	0	0	0	315	78.8	4
WHALE COVE	1982	0	0	0	0	71	29	0	0	0	0	0	0	100	50.0	2
	1984	0	0	0	0	11	13	0	0	0	0	0	0	24	12.0	2
	1985	0	0	0	0	68	12	19	0	5	0	0	0	104	26.0	4
KEEWATIN	1982	0	0	0	0	196	1186	407	203	2	0	0	9	2003	333.8	6
	1983	0	0	0	0	92	466	0	9	19	0	0	0	586	146.5	4
	1984	0	0	0	0	647	834	0	32	9	0	0	0	1522	380.5	4
	1985	0	0	0	0	762	513	22	1	8	0	0	0	1306	261.2	5

For footnotes see Appendix 4.

Table 52. The estimated harvest of snow geese by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
BAKER LAKE	1984	0	0[2]	0	0	149	201	0	0	0	0	0	0	350	175.0[3]	2
	1985	0	0	0	0	0	30	0	0	0	0	0	0	30	30.0	1
CORAL HARBOUR	1982	.[1]	0	.	.	.	4345	0	0	21	0	0	22	4388	1462.7	3
	1983	0	0	0	0	.	.	.	203	12	0	0	0	215	107.5	2
	1984	3	21	0	5	322	5063	0	30	113	0	0	0	5557	793.9	7
	1985	0	0	0	0	118	3798	24	301	12	0	0	0	4253	850.6	5
CHESTERFIELD INLET	1981	.	.	.	.	.	.	.	.	.	0	0	0	0	0.0	0
	1982	0	0	.	.	.	.	.	13	7	19	0	0	39	13.0	3
	1985	0	0	0	0	0	0	0	0	9	0	0	0	9	9.0	1
ESKIMO POINT	1982	0	0	0	0	667	38	0	0	9	0	0	0	714	238.0	3
	1983	0	0	0	0	0	90	0	0	3	0	0	0	93	46.5	2
	1984	0	0	0	0	108	14	0	0	1	0	0	0	123	41.0	3
	1985	0	0	0	0	518	187	0	0	0	0	0	0	705	352.5	2
REPULSE BAY	1982	0	0	0	0	19	8	0	0	0	0	0	.	27	13.5	2
	1984	0	0	0	0	4	0	0	0	0	0	0	0	4	4.0	1
RANKIN INLET	1982	0	0	0	9	0	0	2	0	41	0	0	0	52	17.3	3
	1983	0	0	0	0	0	80	0	0	18	0	0	0	98	49.0	2
	1984	0	0	0	0	209	51	0	0	41	0	0	0	301	100.3	3
	1985	0	0	0	0	184	335	0	0	0	0	0	0	519	259.5	2
WHALE COVE	1982	0	0	0	0	109	40	0	0	0	0	0	0	149	74.5	2
	1984	0	0	0	0	226	273	37	0	4	0	0	0	540	135.0	4
	1985	0	0	0	0	55	0	0	0	29	0	0	0	84	42.0	2
KEEWATIN	1981	.	.	.	.	.	.	.	.	.	0	0	0	0	0.0	0
	1982	0	0	0	9	795	4431	2	13	78	19	0	22	5369	671.1	8
	1983	0	0	0	0	0	170	0	203	33	0	0	0	406	135.3	3
	1984	3	21	0	5	1018	5602	37	30	159	0	0	0	6875	859.4	8
	1985	0	0	0	0	875	4350	24	301	50	0	0	0	5600	1120.0	5

For footnotes see Appendix 4.

Table 53. The estimated harvest of charr by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
BAKER LAKE	1981	.	[1]	.	.	.	.	.	.	.	.	.	[2]	.	[3]	
	1984	0	0	0	0	0	138	0	0	65	0	128	0	128	128.0	1
	1985	0	0	0	0	100	57	0	0	0	4	42	2	247	61.8	4
CORAL HARBOUR	1982	.	318	.	.	.	1504	882	1546	26	12	0	7	168	42.0	4
	1983	0	0	0	0	.	.	.	1066	140	616	53	0	4341	620.1	7
	1984	300	12	2	210	108	370	278	690	29	1110	366	57	2245	449.0	5
	1985	0	202	331	0	138	1093	867	817	11	350	140	484	3733	311.1	12
	1986	27	40	23	.	.	.	.	.	.	.	1281	182	5272	527.2	10
CHESTERFIELD INLET	1982	76	0	.	.	.	.	.	0	0	0	.	.	90	30.0	3
	1983	0	0	0	0	0	55	43	55	0	0	0	0	76	76.0	1
	1984	0	0	0	0	1	0	15	414	50	0	0	0	153	51.0	3
	1985	0	0	0	0	0	8	0	0	0	0	0	0	480	120.0	4
	1986	0	0	13	.	.	.	.	.	.	.	70	0	78	39.0	2
ESKIMO POINT	1981	.	.	.	.	.	.	.	.	.	.	.	.	13	13.0	1
	1982	4	6	0	0	1	72	1080	1042	169	47	24	34	105	35.0	3
	1983	3	0	0	0	61	364	562	694	258	36	57	12	2479	247.9	10
	1984	0	0	0	2	163	172	1249	608	152	85	38	30	2095	232.8	9
	1985	1	0	1	4	41	289	659	1779	357	78	109	68	2601	289.0	9
	1986	37	0	0	.	.	.	.	.	.	59	119	93	3402	309.3	11
REPULSE BAY	1981	.	.	.	.	.	.	.	.	.	.	.	.	37	37.0	1
	1982	0	0	23	0	157	313	47	390	29	133	289	283	705	235.0	3
	1983	0	0	14	0	0	63	76	109	2	542	418	.	1919	239.9	8
	1984	0	0	0	0	14	655	265	493	9	147	564	52	1027	128.4	8
	1985	260	0	0	30	195	501	1017	749	0	631	426	350	2843	355.4	8
	1986	0	0	63	.	.	.	.	.	.	27	639	0	3418	427.3	8
RANKIN INLET	1981	.	.	.	.	.	.	.	.	.	.	.	.	63	63.0	1
	1982	0	353	0	134	82	913	7366	1386	115	791	0	719	719	719.0	1
	1983	214	0	0	0	0	420	788	2117	65	102	727	359	12226	1222.6	10
	1984	333	86	93	57	74	861	724	1804	45	70	443	492	4641	580.1	8
	1985	124	44	96	114	139	1290	1448	2873	0	404	763	205	5115	426.3	12
	1986	315	114	107	.	.	.	.	.	.	.	615	316	7463	678.5	11
WHALE COVE	1981	.	.	.	.	.	.	.	.	.	.	.	.	536	178.7	3
	1982	111	42	79	58	86	332	630	6583	44	42	11	166	219	73.0	3
	1983	2	25	0	.	.	.	.	.	.	36	69	13	8083	673.6	12
	1984	23	0	0	7	3	88	261	467	20	195	63	30	120	30.0	4
	1985	4	5	0	0	56	76	117	416	3	10	117	16	1197	119.7	10
	1986	40	20	3	.	.	.	.	.	.	.	9	6	702	70.2	10
KEEWATIN	1981	.	.	.	.	.	.	.	.	.	.	.	.	63	21.0	3
	1982	191	719	102	192	326	3134	10005	10947	383	222	452	1202	1876	625.3	3
	1983	219	25	14	0	61	902	1469	4041	465	1417	1324	384	29124	2427.0	12
	1984	656	98	95	276	363	2284	2792	4476	370	950	1474	661	10281	934.6	11
	1985	389	251	428	148	669	3314	4108	6634	371	2084	1597	1125	16216	1351.3	12
	1986	419	174	209	.	.	.	.	.	.	854	2733	604	20503	1708.6	12
											.	.	.	802	267.3	3

For footnotes see Appendix 4.

Table 54. The estimated harvest of lake trout by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	$\bar{X}$	NUM
BAKER LAKE	1981	[1] .	.	.	.	.	.	.	.	.	.	8744	2154	10898	5449.0[3]	2
	1982	0	0	380	400	0	0	0	.	0	.	0	0	780	390.0	2
	1983	0	0	0	0	796	0	1886	188	0	1732	545	178	5325	887.5	6
	1984	76	157	268	241	32	175	72	87	182	196	144	192	1822	151.8	12
	1985	181	43	0	0	264	596	333	59	25	798	520	469	3288	328.8	10
	1986	261	317	250	.	.	.	.	.	.	.	.	.	828	276.0	3
CORAL HARBOUR	1982	.	[2] 0	.	.	.	0	0	0	0	0	0	9	9	9.0	1
	1983	0	0	410	0	.	.	.	0	0	0	0	0	410	410.0	1
	1985	0	0	0	0	0	7	0	0	0	0	0	0	7	7.0	1
CHESTERFIELD INLET	1982	0	19	.	.	.	.	.	202	0	26	26	0	273	68.3	4
	1983	0	0	30	131	41	72	8	0	0	0	43	0	325	54.2	6
	1984	0	0	0	1	59	26	0	0	0	111	0	0	197	49.3	4
	1985	0	0	0	2	31	22	0	0	0	0	58	0	113	28.3	4
ESKIMO POINT	1981	.	.	.	.	.	.	.	.	.	879	679	128	1686	562.0	3
	1982	91	12	79	182	237	38	9	65	75	28	5	0	821	74.6	11
	1983	48	11	213	115	281	72	0	24	130	68	159	231	1352	122.9	11
	1984	0	0	4	183	62	138	8	6	111	22	22	32	588	58.8	10
	1985	4	0	0	0	0	0	0	0	0	75	59	32	170	42.5	4
	1986	2	4	70	.	.	.	.	.	.	.	.	.	76	25.3	3
REPULSE BAY	1981	.	.	.	.	.	.	.	.	.	745	5	0	750	375.0	2
	1982	75	0	1	15	199	345	0	0	11	0	5	.	651	93.0	7
	1983	0	0	8	46	8	2	0	0	0	2	0	0	66	13.2	5
	1984	0	0	0	0	60	0	0	0	0	0	225	0	285	142.5	2
	1985	0	0	0	0	68	115	8	0	0	16	0	1	208	41.6	5
	1986	0	0	4	.	.	.	.	.	.	.	.	.	4	4.0	1
RANKIN INLET	1981	.	.	.	.	.	.	.	.	.	.	0	33	33	33.0	1
	1982	0	15	78	59	0	0	0	0	0	0	45	0	197	49.3	4
	1983	0	20	0	0	289	0	0	0	0	0	54	0	363	121.0	3
	1984	0	20	0	120	232	0	10	21	0	0	103	0	506	84.3	6
	1985	1	16	14	0	173	127	0	0	15	0	0	2	348	49.7	7
	1986	16	0	0	.	.	.	.	.	.	.	.	.	16	16.0	1
WHALE COVE	1981	.	.	.	.	.	.	.	.	.	4	33	156	193	64.3	3
	1982	59	12	14	71	52	15	23	120	3	15	26	58	468	39.0	12
	1983	11	56	18	.	.	.	.	.	.	.	0	0	85	28.3	3
	1984	17	14	92	93	80	18	0	0	0	0	9	0	323	46.1	7
	1985	0	0	0	6	52	44	7	1	0	3	0	0	113	18.8	6
	1986	10	9	0	.	.	.	.	.	.	.	.	.	19	9.5	2
KEEWATIN	1981	.	.	.	.	.	.	.	.	.	1628	9461	2471	13560	4520.0	3
	1982	225	58	552	727	488	398	32	387	89	69	107	67	3199	266.6	12
	1983	59	87	679	292	1415	146	1894	212	130	1802	801	409	7926	660.5	12
	1984	93	191	364	638	525	357	90	114	293	329	503	224	3721	310.1	12
	1985	186	59	14	8	588	911	348	60	40	892	637	504	4247	353.9	12
	1986	289	330	324	.	.	.	.	.	.	.	.	.	943	314.3	3

For footnotes see Appendix 4.



Fig. 1. Map of Keewatin District showing the seven communities surveyed during the harvest study and the zonal grid used to locate kills.

Appendix 1. Monthly percent of hunters interviewed for the seven Keewatin communities for the period October 1981 to March 1986.

	Year	Oct.	Nov.	Dec.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.
Baker Lake	1981-82	*[1]	27.3	18.6	45.5	56.4	49.5	40.9	57.3	39.5	41.8	*	90.9
	1982-83	*	94.1	94.5	99.6	85.5	91.8	95.7	93.2	94.5	92.3	95.9	92.7
	1983-84	98.7[2]	93.2	97.8	96.1	96.7	95.2	97.4	94.4	97.0	95.9	96.6	95.8
	1984-85	97.6	97.7	98.0	96.8	96.6	94.9	96.5	100.0	98.6	99.0	95.0	96.7
	1985-86	95.4	98.0	94.8	100.0	100.0	100.0						
Chesterfield Inlet	1981-82	*	*	*	68.0	48.0	*	*	*	*	*	24.0	88.0
	1982-83	78.0	86.0	90.0	62.0	56.0	28.0	58.0	88.0	100.0	90.0	100.0	100.0
	1983-84	100.0	88.3	98.3	93.3	92.7	100.0	100.0	78.1	96.9	87.0	98.4	100.0
	1984-85	100.0	100.0	100.0	100.0	100.0	98.5	98.3	95.8	95.7	97.2	93.2	97 1
	1985-86	100.0	100.0	100.0	100.0	92.0	91.8						
Coral Harbour	1981-82	*	*	*	*	49.5	*	*	*	56.2	59.0	36.2	73.3
	1982-83	18.1	27.6	26.7	31.4	32.4	35.2	18.1	*	*	*	*	*
	1983-84	27.6	22.9	16.2	32.4	95.2	85.7	82.9	69.5	96.3	73.3	67.6	60.0
	1984-85	99.1	61.9	62.9	68.6	79.0	93.3	94.3	78.1	59.0	62.9	75.2	86.7
	1985-86	73.3	85.7	80.0	85.0	84.8	81.0						
Eskimo Point	1981-82	57.5	62.8	61.1	79.6	78.3	66.4	50.0	81.4	79.2	80.5	80.1	88.2
	1982-83	97.3	94.7	95.6	97.8	97.0	100.0	97.5	96.7	95.5	96.2	98.4	97.7
	1983-84	98.8	98.4	97.0	98.3	94.6	93.2	93.9	99.1	98.2	99.2	98.4	84.1
	1984-85	86.3	94.5	79.2	93.8	89.3	90.1	95.2	88.3	89.4	82.3	95.2	94.3
	1985-86	87.5	91.9	91.8	94.1	85.8	89.8						
Rankin Inlet	1981-82	*	92.3	57.5	92.7	66.8	71.0	33.7	31.6	45.6	63.7	22.8	96.9
	1982-83	61.7	64.8	100.0	84.5	89.1	84.5	95.9	33.7	19.2	34.2	55.4	61.1
	1983-84	51.3	81.5	92.2	90.4	89.5	91.0	97.5	74.4	90.3	82.4	100.0	85.8
	1984-85	84.7	82.1	86.0	86.7	85.4	91.4	94.8	87.6	87.3	88.6	93.5	96.2
	1985-86	95.0	98.1	97.5	97.9	96.8	97.2						
Repulse Bay	1981-82	61.1	58.9	26.7	38.9	60.0	26.7	50.0	31.1	37.8	32.2	33.3	44.4
	1982-83	51.1	56.7	*	70.0	73.3	71.1	*	53.3	53.3	73.3	57.8	53.3
	1983-84	57.8	66.7	41.1	58.9	44.4	54.4	51.1	73.3	71.1	51.1	76.7	58.9
	1984-85	67.9	72.6	70.2	79.8	66.7	73.8	73.8	67.9	61.9	72.6	73.6	75.0
	1985-86	71.4	83.3	97.7	90.3	95.2	100.0						
Whale Cove	1981-82	20.0	86.0	82.0	82.0	74.0	94.0	76.0	52.0	100.0	100.0	72.0	92.0
	1982-83	28.0	32.0	54.0	62.0	80.0	40.0	*	*	*	*	*	*
	1983-84	*	30.0	14.0	52.0	98.0	98.0	100.0	77.6	70.7	69.9	71.0	88.7
	1984-85	*	68.0	85.9	87.5	93.5	93.3	87.3	83 9	82.5	98.3	98.3	100.0
	1985-86	100.0	100.0	100.0	98.3	100.0	100.0						

[1]Data not provided.
[2]These figures are the percentage of hunters in each community who were interviewed each month in relation to the total number of hunters who could have hunted that month.

Appendix 2. Monthly theoretical kill factors for seven Keewatin communities for the period October 1981 to March 1986.

	Year	Oct.	Nov.	Dec.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.
Baker Lake	1981-82	*[3]	(3.67)[1]	(5.36)[4]	2.19	1.77	2.02	2.44	1.75	2.52	2.38	*	1.11
	1982-83	*	1.06	1.06	1.00	1.19	1.08	1.00	1.01	1.03	1.03	1.04	1.06
	1983-84	1.00	1.07	1.00	1.00	1.00	1.00	1.00	1.08	1.00	1.00	1.00	1.00
	1984-85	1.00	1.00	1.00	1.00	1.00	1.00	1.01	1.00	1.01	1.00	1.00	1.00
	1985-86	1.00	1.00	1.00	1.00	1.00	1.00						
Chesterfield Inlet	1981-82	*	*	*	1.47	2.08	*	*	*	*	*	(4.17)	1.13
	1982-83	1.28	1.16	1.11	1.61	1.78	(3.57)	1.72	1.14	1.00	1.18	1.00	1.00
	1983-84	1.00	1.00	1.04	1.14	1.27	1.00	1.00	1.35	1.07	1.39	1.03	1.00
	1984-85	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.13	1.11	1.07	1.21	1.08
	1985-86	1.00	1.00	1.00	1.00	1.00	1.00						
Coral Harbour	1981-82	*	*	*	*	2.01	*	*	*	1.78	1.69	2.76	1.36
	1982-83	(5.53)	(3.62)	(3.75)	3.17	3.08	2.83	5.53	*	*	*	*	*
	1983-84	(3.62)	(4.38)	(6.18)	(3.09)	1.05	1.17	1.21	1.44	1.01	1.35	1.44	1.61
	1984-85	1.01	1.54	1.40	1.22	1.59	1.04	1.02	1.28	1.65	1.52	1.29	1.09
	1985-86	1.62	1.27	1.57	1.00	1.10	1.07						
Eskimo Point	1981-82	1.74	1.59	1.64	1.25	1.28	1.51	2.00	1.23	1.18	1.24	1.23	1.14
	1982-83	1.03	1.06	1.02	1.07	1.11	1.11	1.05	1.04	1.08	1.06	1.03	1.05
	1983-84	1.38	1.01	1.00	1.00	1.05	1.01	1.01	1.01	1.02	1.01	1.02	1.12
	1984-85	1.09	1.01	1.22	1.04	1.14	1.31	1.17	1.14	1.11	1.40	1.07	1.01
	1985-86	1.07	1.21	1.24	1.13	1.48	1.21						
Rankin Inlet	1981-82	*	1.08	1.74	1.05	1.50	1.41	2.97	3.16	2.19	1.56	(4.39)	1.03
	1982-83	1.62	1.54	1.00	1.18	1.06	1.17	1.04	2.95	(5.22)	2.92	1.80	1.63
	1983-84	1.95	1.15	1.22	1.16	1.18	1.02	1.06	1.42	1.00	1.38	1.00	1.06
	1984-85	1.18	1.36	1.36	1.33	1.63	1.37	1.35	1.39	1.30	1.96	1.25	1.16
	1985-86	1.05	1.05	1.09	1.06	1.10	1.08						
Repulse Bay	1981-82	1.64	1.70	(3.75)	2.57	1.66	(3.75)	1.70	3.21	2.65	3.10	3.00	2.25
	1982-83	1.95	1.70	*	1.43	1.36	1.40	*	1.52	1.67	1.36	1.73	1.87
	1983-84	1.78	1.42	2.36	1.66	2.04	1.76	1.76	1.36	1.25	1.80	1.27	1.55
	1984-85	1.38[2]	1.34	1.40	1.20	1.38	1.27	1.31	1.39	1.51	1.35	1.33	1.31
	1985-86	1.33	1.16	1.07	1.23	1.10	1.00						
Whale Cove	1981-82	(5.00	1.16	1.21	1.21	1.35	1.06	1.31	1.92	1.00	11.00	1.39	1.09
	1982-83	(3.57)	(3.13)	1.85	1.61	1.25	2.50	*	*	*	*	*	*
	1983-84	*	(3.33)	(7.14)	7.92	1.01	1.02	1.00	1.13	1.47	1.48	1.45	1.07
	1984-85	*	1.32	1.04	1.00	1.00	1.03	1.00	1.06	1.05	1.00	1.00	1.00
	1985-86	1.00	1.00	1.00	1.00	1.00	1.00						

[1]Bracketed figures were not used because they were based on insufficient data.
[2]The figure for known hunters in Repulse Bay was lowered to 84 from 90 beginning October, 1985. This figure is important when determining the number of hunters not contacted on a given month with in a community.
[3]*Data not provided.
( )Data provided on successful hunters only. Estimated harvest = Reported harvest.

Appendix 3. Members of the Steering Committee for the period of the Keewatin Wildlife Federation Harvest Study, 1981 to 1987.

Chairperson

Mr. F. McFarland and Ms. D. Stewart	Northern Affairs Program, Department of Indian Affairs and Northern Development.	1981-85 1985-87

Members

Mr. R. Bailey Mr. R. Cole	Canadian Wildlife Service, Department of the Environment.	1981-82 1982-87
Mr. K. Lloyd Mr. R. Graf	Department of Renewable Resources, Government of the Northwest Territories.	1981-82 1982-87
Mr. R. Peet	Department of Fisheries and Oceans.	1981-87
Mr. L. Pilikapsi	President, Keewatin Wildlife Federation.	1981-82
Mr. P. Kritterdiluk	President, Keewatin Wildlife Federation.	1982-83
Mr. D. Milortok	President, Keewatin Wildlife Federation.	1983-85
Mr. A. Anyootealuk	President, Keewatin Wildlife Federation.	1985-86
Mr. J. Kaunak	President, Keewatin Wildlife Federation.	1986-87
Mr. L. Gamble	Regional Resource Manager, Keewatin Harvest Study.	1981-87
Mr. L. Guluk	Project Manager, Keewatin Harvest Study.	1981-83
Ms. V. Curley	Assistant Regional Resource Manager, Keewatin Harvest Study.	1983-85
Mr. J. Kusugak	Assistant Regional Resource Manager, Keewatin Harvest Study.	1985-86

Appendix 4. Footnotes for Tables 35 to 54.

A) Footnotes for Tables 35 to 41.

[1] In the case where there was zero harvest for all months this row is not represented in the table but is reflected in the N row.

[2] Zero means there was no harvest for that particular month.

[3] A dot means the data were not available for that month. When this occurred that month is not counted in the row N for that year or in the column N for that month.

[4] In cases where there was minimal harvest column N, sums and means have not been included to conserve space.

[5] Categories are as follows: M = males, F = females, C = calf, and U = unknown.

[6] Bracketted letters indicate the following: H = high profile species, L = low profile species, A = abundant, S = scarce. These categories are defined as follows:

1. A high profile species is one which is of economic and cultural importance in a particular community. This may vary between communities because the availability for harvest influences the importance of specific species.
2. A low profile species is one which has little economic or cultural significance no matter what its abundance.
3. Abundant refers primarily to availability for harvest. This may vary with distance or with time. For example some species, such as caribou may be available to a community year-round but their harvest may involve considerable travel at certain periods of the season. Other species such as beluga whale and geese are only abundant at specific locations and periods in each year.
4. Scarce indicates the availability of a species for harvest is limited or that the effort involved in harvesting it is too great. In many instances the harvest of such species is incidental to more preferred species which are actively sought.

B) Footnotes for Tables 42 to 54.

[1] A dot means the data were not available for that month.

[2] Zero means there was no harvest for that particular month.

[3] $\bar{X}$ was calculated using only those months showing a harvest.

Printed in Canada by
Supply and Services Canada
Printing Office
for exclusive distribution by
Fisheries and Oceans
Freshwater Institute
Winnipeg, Manitoba
Canada